



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

UCRL-CONF-202594

Very Fine Aerosols from the World Trade Center Collapse Piles: Anaerobic Incineration?

***T.A. Cahill, S.S. Cliff, J. Shackelford,
M. Meier, M. Dunlap, K.D. Perry,
G. Bench, and R. Leifer***

September, 12, 2003

American Chemical Society Meeting, Sep 7-12, 2003, New York,
NY

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

RESERVE THIS SPACE

Very fine aerosols from the World Trade Center collapse piles: Anaerobic incineration?

**Thomas A. Cahill¹, Steven S. Cliff¹, James Shackelford¹,
Michael Meier¹, Michael Dunlap¹, Kevin D. Perry², Graham
Bench³, and Robert Leifer⁴**

**¹The D. E. L. T. A. Group, Depts. of Applied Science and Chemical
Engineering/Materials Science, University of California, Davis
CA 95616**

²Dept. of Meteorology, University of Utah

**³Center for Accelerator Mass Spectrometry, Lawrence
Livermore National Laboratory**

**⁴U.S. Department of Homeland Security, Environmental
Measurements Laboratory, 201 Varick Street, New York**

By September 14, three days after the initial World Trade Center collapse, efforts at fire suppression and heavy rainfall had extinguished the immediate surface fires. From then until roughly mid-December, the collapse piles continuously emitted an acrid smoke and fume in the smoldering phase of the event. Knowledge of the sources, nature, and concentration of these aerosols is important for evaluation and alleviation of the health effects on workers and nearby residents. In this paper, we build on our earlier work to ascribe these aerosols to similar processes that occur in urban incinerators. The simultaneous presence of finely powdered (circa 5 μm) and highly basic (pH 11 to 12) cement dust and high levels of very fine (<0.25 μm) sulfuric acid fumes help explain observed health impacts. The unprecedented levels of several

RESERVE THIS SPACE

metals in the very fine mode can be tied to liberation of those metals that are both present in elevated concentrations in the debris and have depressed volatility temperatures caused by the presence of organic materials and chlorine.

Introduction

The collapse of the World Trade Center structures (South Tower, North Tower, and WTC 7) presented two very different types of air pollution events:

1. Initial fires and collapse - derived "dust storm"
2. Continuing emissions from the debris piles

Both cases shared the unusual aspect of a massive ground level source of particulate matter in a highly populated area with potential health impacts. EPA (Devlin, 2002) summarized 5 causal factors most likely to explain the statistically solid data connecting fine PM_{2.5} aerosols and human health.

1. Biological aerosols (bacteria, molds, viruses...)
2. acid aerosols
3. very fine/ultrafine (<0.1 μm) insoluble aerosols
4. fine transition metals
5. high temperature organics

We especially needed to ascertain if these potentially causal factors were present in the WTC plumes.

The presence of fuels, including diesel and electrical insulating oils and combustible materials in the WTC buildings, and lack of oxygen were two factors that allowed the fire to burn for over 3 months. In order to better understand these aerosols, a DELTA Group rotating DRUM impactor was shipped to New York and set upon the roof of the DOE Environmental Measurement Laboratory (EML) at 201 Varick Street, 45 m above ground level and roughly 1.5 km NNE of the World Trade Center site. The DRUM operated until late December, after the last surface fires had been extinguished. (Cahill et al, 2004)

The samples were collected in 8 size modes from the inlet (circa 12 μm) to 5.0, 2.5, 1.15, 0.75, 0.56, 0.34, 0.26 to 0.09 μm aerodynamic diameter (Cahill et al, 1985; Raabe et al, 1989; Raabe 1997). Samples were analyzed by a suite of non-destructive spatially resolved beam based techniques (Bench et al, 2002; DQAP v.9/02, 2002) for mass, hydrogen as an organics surrogate, elements from sodium to molybdenum and some heavy elements by synchrotron x-ray fluorescence (S-XRF), particle size by scanning electron microscopy. Time resolution ranges from 1½ hr to 3 hr depending on the size of the exciting beam.

The presence of the WTC plume at Varick Street depended on both source emission rate and meteorology, with both wind direction and plume lofting as the key parameters. In addition, the nature of the aerosols themselves, especially as they deviated in concentration, size, and composition from upwind and regional aerosols, also provided information identifying WTC impacts. We chose the 5 parameters that deviated generally by an order of magnitude from typical regional and urban values:

1. Very fine ($0.26 > D_p > 0.09 \mu\text{m}$) aerosol mass $> 3.0 \mu\text{g}/\text{m}^3$, ($> 10 \times$ background), in 3 to 6 hr plumes.

18 such events, typically 3 to 6 hr in duration at Varick Street, were seen in the month of October. Then we applied the additional criteria:

- | | #events |
|--|---------|
| 2. HYSPLIT wind in SSW quadrant 14+1 calm | |
| 3. Very fine organics $> 1.0 \mu\text{g}/\text{m}^3$, | 16 |
| 4. 2.5 to 5 μm cement dust EF > 2.5 | 16 |
| 5. 2.5 to 5 μm sulfate $> 0.3 \mu\text{g}/\text{m}^3$ | 15 |
| 6. Simultaneous ground based 3 hr haze plumes at La Guardia airport, $L_v < 15 \text{ km}$ 5 (+3 days Sept.) | |

5 events met all 6 criteria, 4 more met 5 of the 6 criteria, and these 9 events are labeled “highly probable events”. Three events met only one or two of the parameters, and these are labeled “non-WTC plumes” or uncertain sources, including a local power plant. Six of the events met 3 or 4 of the parameters, and these were labeled “probable WTC influence”. The analysis that follows uses only data from the 9 highly probable events.

On most days, the plume sloped above NYC so that only those on or near the WTC site breathed these aerosols.

Table 6 shows very fine particle values for six of the highly probable events plus background, a non-WTC plume event, and comparison data from two highly impacted sites using the same size and plume behavior from earlier studies. (Cahill et al, 1992; Reid et al, 1994; Seinfeld et al, 2004).

Date	WTC impact?	Mass ($\mu\text{g}/\text{m}^3$)	Organics ($\mu\text{g}/\text{m}^3$)	SiO_2 ($\mu\text{g}/\text{m}^3$)	H_2SO_4 ($\mu\text{g}/\text{m}^3$)	V (ng/m^3)	Ni (ng/m^3)
Oct.7	No	0.5	0.04	0.02	0.1	0.1	0.1
Oct.3	Yes	50.6	9.3	1.4	17.1	115	22.7
Oct.4	Yes	9.3	3.2	2.6	3.7	61	8.0
Oct.5	Yes	4.7	1.5	0.9	1.9	3	1.3
Oct.12	Yes	8.2	3.0	2.9	2.2	24	7.1
Oct.15	Yes	4.8	1.6	1.3	1.1	3	17.3
Oct.24	Yes	3.8	1.0	0.2	1.6	5	6.7
Oct.29	No	2.4	1.2	0.07	0.9	2	1.6
Kuwait	-	na	na	0.6	5.5	na	5.0

Beijing	-	na	na	1.1	6.7	0.8	1.8
---------	---	----	----	-----	-----	-----	-----

Table 6 Results of aerosol sampling at Varick Street, October, 2001, for very fine ($0.26 > D_p > 0.09 \mu\text{m}$) particles.

Coarse particles were similar to the initial collapse aerosols (cement, drywall, glass, ...) (Lioy et al., 2003), but had chemicals and soot from the ongoing combustion (Natusch et al., 1974a,b). The presence of an unprecedented (versus Beijing, Kuwait) level of very fine ($0.26 > D_p > 0.09 \mu\text{m}$) particles by mass and number in narrow plumes was more typical of an industrial source. Upwind sources were a very minor contribution as shown by direct comparison with upwind aerosol sites and size distributions grossly different from typical ambient aerosols (Leifer et al., this volume). Very fine particles near the WTC site in May, 2002, were generally <10% of the October, 2001 plume impact days at Varick Street.

Interpretation

Figure 1 shows an example of typical crustal elements (silicon, calcium, iron, and aluminum) in the very fine ($0.26 > D_p > 0.09 \mu\text{m}$) mode from October 2 to October 30, 2001.




Figure 1 Typical crustal elements (silicon, calcium, iron, and aluminum) in the very fine particle ($0.26 > D_p > 0.09 \mu\text{m}$) mode from October 2 through October 30, 2001. On the top is shown the reflected light picture of this DRUM stage, and below the plot the meteorological favored wind direction that would bring the plume to Varick Street. Two rain events and an easterly ocean wind are also shown.

There are several important points to note in this figure. First, the other crustal elements (Al, Ca, and iron) seen so abundantly in the coarser particles are essentially absent in this very fine mode. Thus, the signal is not some sampling artifact, as is also shown by the low silicon values in the sub-micron sizes, and the mechanisms are not simply mechanical grinding. Second, the silicon only appears when the wind is correct. Third, the peak values trend downward in October, and even the color grows fainter. October 24 was by all accounts (meteorology, coarse calcium, ...) a very strong plume impact, but by then the silicon has gone. The same pattern is seen

in some other anthropogenic elements, notably vanadium), in both the coarse and fine modes.

These data then pose problems of the aerosol sources. We see aerosol typical of a high temperature industrial source, yet the temperatures in the near sub-surface collapse piles were not very high. We see some elements vastly enhanced over typical Earth crustal values and others, equally present in the dust, absent in the aerosols. We propose a model derived from the theory and data on municipal incinerators. It was discovered that the presence of organic matter and especially chlorine in the wastes liberated metals by greatly reducing their volatility temperatures. (Barton et al, 1990; Seeker et al, 1990, and references therein).

In Table 2, we present these data for EPA criterion metals and a few others of interest. The data are ordered by volatility temperature (the point at which the vapor pressure reaches 10^{-6} atmospheres).

Incineration with 10% chlorine	Boiling Point	Earth Crustal	Bulk Dust EPA	Bulk dust Liroy	Volatility Temp	Principal Species
Metal	°C	ppm	ppm	ppm	°C	
Chromium	2639	102	71.5	165	1594	CrO ₂ /O ₃
Beryllium	1280	2.8	1.7	3.2	1042	Be(OH) ₂
Barium	1634	425	195	381	895	BaCl ₂
Nickel	2834	84	15.5	43.5	686	NiCl ₂
Antimony	697	0.2	na	na	653	Sb ₂ O ₃
Silver	2190	0.004	4.9	2.3	620	AgCl
Selenium	Na	0.05	<1	na	315	SeO ₂
Cadmium	761	0.15	3.8	7.2	211	Cd
Vanadium	3480	120	18.3	38.9	147	VCl ₄
Thallium	1464	9.6	<1	1.4	136	TlOH
Osmium	4224	0.0015	Na	na	40	OsO ₄
Arsenic	814	1.8	<1	2.6	32	As ₂ O ₃

Mercury	353	0.085	0.37	nd	25	Hg
SiO ₂	1725	9000	na	na	12	SiCl ₄
Lead	1748	14	98	305	-12	PbCl ₄

Table 2 Concentration in the WTC dusts and volatility temperatures of elements in the presence of 10% chlorine.

Note that the level of many metals is greatly enhanced over Earth crustal levels, as to be expected in the wreckage of a highly computerized building. The top elements, chromium, beryllium, and barium, all have volatility temperatures that are probably higher than most parts of the collapse piles, since steel was not melted but was observed to glow red when pulled from the piles. Nickel, antimony, and silver are marginal, but all the rest have volatility temperatures well below the documented temperature of the collapse piles during October.

The mechanism proposed is then the formation of gasses in the oxygen-poor, chlorine-rich collapse piles, emission through the rubble and conversion at or near the surface into oxidized forms and thus very fine particles.

The other factor is concentration in the WTC dusts. The elements beryllium (which we cannot analyze by S-XRF), antimony, silver, selenium, cadmium, thallium, osmium, arsenic and mercury are present only at the ppm levels in the WTC dusts, and thus would be unlikely in aerosols no matter what was the volatility temperature.


Combining these data into table 3, we see that the elements actually observed share the characteristics of relatively high concentrations in the collapse piles and low volatility temperatures. Other species, including most other crustal elements absent in the very fine mode, do not have depressed volatility temperatures and thus are not seen in the very fine aerosol. Chromium and barium ,

both with high volatility temperatures, although abundant in the dusts, are largely absent in the aerosols. Vanadium and nickel are equally present in the dust, but because of the much lower vanadium volatility temperature, vanadium has 5 times more mass in the aerosol than nickel.

Very fine aerosols	October 7 background	October 3 WTC plume	Average abundance	Volatility temperature
0.26–0.09 μm	$\mu\text{g}/\text{m}^3$	$\mu\text{g}/\text{m}^3$	WTC dust	10% chlorine
Mass	0.53	50.7	na	na
Organics	0.04	9.3	na	na
Sulfur	0.04	5.6	na	na
	ng/m^3	ng/m^3	ppm	$^{\circ}\text{C}$
Silicon	11	698	abundant	12
Vanadium	0.1	114	30	147
Lead	<0.5	26	200	-12
Nickel	0.1	23	30	686
Chromium	<0.1	1.5	120	1594
Barium	<0.1	<0.5	290	895

Table 3 Very fine aerosols versus concentration in dusts and volatility temperature.

Further evidence for the role of chlorine in releasing certain metals from the collapse pile is shown in Figure 2. Figure 2 shows vanadium and chlorine in the very fine mode aerosols but at a ratio well below that of VCl_4 .



Conclusions

By September 14, three days after the initial World Trade Center collapse, efforts at fire suppression and heavy rainfall had extinguished the immediate surface fires. From then until roughly mid-December, the collapse piles continuously emitted an acid smoke and fume in the smoldering phase of the event. Knowledge of the sources, nature, and concentration of these aerosols is important for evaluation and alleviation of the health effects on workers and nearby residents. The simultaneous presence of very fine powdered and highly basic (pH 11 to 12) cement dust and high levels of very fine ($<0.25 \mu\text{m}$) sulfuric acid fumes help explain observed health impacts. The unprecedented levels of several metals in the very fine mode can be tied to liberation of those metals that are both present in the debris and have depressed

volatility temperatures caused by the presence of organic materials and chlorine. These plumes have the potential for health impacts, as potentially causal factors if health in very fine particles reached unprecedented ambient levels in the very fine aerosol plumes from the WTC collapse piles. This would affect workers at the site who did not use adequate respirators but because of plume lofting would not have affected most New York residents away from the site.

Acknowledgements

The authors would like to gratefully acknowledge the scores of people and numerous organizations that contributed their time and effort to this work, all of which was done on a volunteer basis. The work benefited greatly from sampler and analytical developments occasioned by the ACE-Asia study NSF ATM 0080225, support from the Center for Accelerator Mass Spectrometry, LLNL, and resources Center for X-ray Optics of the Advanced Light Source, LBNL. This work was performed in part under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

We would like to especially thank William Wilson, USEPA, for his careful review and thoughtful discussions which led to significant improvements in the paper.

The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model (<http://www.arl.noaa.gov/ready.html>) and the READY website used in this publication.

We gratefully acknowledge the support of the American Lung Association for the May, 2002 study,

We wish to thank all the people who helped increase our understanding of the WTC problem, and in particular the very helpful analysis of Prof. Ian Kennedy (UCD Engineering) and Dr. William Wilson, (ORD, USEPA).

References

- Barton, R. G., W. D. Clark, and W. R. Seeker, Fate of Metals in Waste Combustion Systems, *Combust. Sci. & Tech.*, 74:327-342 (1990)
- Bench, Graham, P. G. Grant, D. Ueda, S. S. Cliff, K. D. Perry, and T. A. Cahill. (2002) The Use of STIM and PESA to Measure Profiles of Aerosol Mass and Hydrogen Content, Respectively, across Mylar Rotating Drums Impactors Samples, *Aerosol Science and Technology* 36, 642-651.
- Cahill, Thomas A. Comments on surface coatings for Lundgren-type impactors. (1979) *Aerosol Measurement*. Dale A. Lundgren, Editor. University Presses of Florida, Pp. 131-134.
- Cahill, T. A., C. Goodart, J. W. Nelson, R. A. Eldred, J. S. Nasstrom, and P. J. Feeney. (1985) Design and evaluation of the drum impactor. *Proceedings of International Symposium on Particulate and Multi-phase Processes*. Teoman Ariman and T. Nejat Veziroglu, Editors. Hemisphere Publishing Corporation, Washington, D.C. 2:319-325.
- Cahill, Thomas A., Marcelle Surovik, and Ian Wittmeyer. (1990) Visibility and aerosols during the 1986 carbonaceous species methods comparison study. *Aerosol Science and Technology*. 12:149-160.
- Cahill, Thomas A., Kent Wilkinson, and Russ Schnell. (1992). Composition analyses of size-resolved aerosol sample taken from

aircraft downwind of Kuwait, Spring, 1991. *Journal of Geophysical Research*. Vol. 97, No. D13, Paper no. 92JD01373, Pp. 14513 -14520, September 20.

Cahill, Thomas A. and Paul Wakabayashi. (1993). Compositional analysis of size-segregated aerosol samples. Chapter in the ACS book *Measurement Challenges in Atmospheric Chemistry*. Leonard Newman, Editor. Chapter 7, Pp. 211 -228.

Cahill, Thomas A. Compositional Analysis of Atmospheric Aerosols. (1995) *Particle-Induced X-Ray Emission Spectrometry*, Edited by Sven A. E. Johansson, John L. Campbell, and Klas G. Malmqvist. *Chemical Analysis Series*, Vol. 133, pp. 237- 311. John Wiley & Sons, Inc.

Cahill, T. A., P. Wakabayashi, T. James. Chemical State of Sulfate at Shenandoah National Park During Summer 1991 *Nuclear Instruments and Methods in Physics Research B: Beam Interactions with Materials and Atoms*, 109/110 (1996) 542 -547.

Devlin, R. "What we know and what we think we know about health effects of fine particles; Is it enough?" AAAR, Charlotte, NC (2003)

DQAP 7.02 UC Davis DELTA Group *DRUM Quality Assurance Protocols*, version 7.02 (2002)

EPA 600/P-95/001af/Vol. 1, 6 -170 to 6 -220 (1996)

HYSPLIT <http://www.arl.noaa.gov/ready/hysplit4.html> (2002)

Lioy, P. J., Weisel, C. P., Millette, J. R., Eisenreich, S., Vallero, D., Offenberg, J., Buckley, B., Turpin, B., Zhong, M., Cohen, M. D., Prophet, C., Yang, I., Stiles, R., Chee, G., Johnson, W., Porcja, R.,

Alimokhtari, S., Hale, R. C., Weschler, C., and Chen, L. C. (2002) Characterization of the Dust/Smoke Aerosol that Settled East of the World Trade Center (WTC) in Lower Manhattan after Collapse of the WTC 11 September, 2001., *Environmental health perspectives* 110, #7, 703–714.

Lundgren, D. A., Journal of the Air Pollution Control Association. 17:225 (1967)

Marple, V., Rubow, K., Anath, G., and Fissan, H. Journal of Aerosol Science 17:489–494 (1986)

Natusch, D. F. S., Wallace, J. R., and Evans, C. A., Science 183:202 (1974)

Natusch, D. F. S., and Wallace, J. R., Science 186:695 (1974)

Raabe, Otto G., David A. Braaten, Richard L. Axelbaum, Stephen V. Teague, and Thomas A. Cahill. (1988) Calibration Studies of the DRUM Impactor. *Journal of Aerosol Science*. 19.2:183-195.

Raabe, O. private communication (1997)

Reid, J. S., T. A. Cahill, E. A. Gearhart, R. G. Flocchini, J. S. Schweitzer, and C. A. Peterson. Elemental analysis of Kuwaiti petroleum and combustion products. (1993) *Journal of Nuclear Geophysics*. Vol. 7, No. 1, Pp. 81–86.