

The Spin-Up Rate of the White Dwarf of GK Per

C.W. Mauche

This article was submitted to
International Astronomical Union Colloquium 190 – Magnetic
Cataclysmic Variables, Cape Town, South Africa, December 8-13,
2002

January 24, 2003

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This report has been reproduced directly from the best available copy.

Available electronically at <http://www.doe.gov/bridge>

Available for a processing fee to U.S. Department of Energy
and its contractors in paper from
U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Facsimile: (865) 576-5728
E-mail: reports@adonis.osti.gov

Available for the sale to the public from
U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Telephone: (800) 553-6847
Facsimile: (703) 605-6900
E-mail: orders@ntis.fedworld.gov
Online ordering: <http://www.ntis.gov/ordering.htm>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

This work was performed under the auspices of the U.S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

The Spin-Up Rate of the White Dwarf of GK Per

Christopher W. Mauche

Lawrence Livermore National Laboratory,
L-473, 7000 East Avenue, Livermore, CA 94550

Abstract. We use hard X-ray light curves measured by the *Chandra* HETG and *RXTE* PCA during the late rise and plateau phases of the 2002 March–April outburst of the intermediate polar GK Per to determine that its X-ray pulse period $P = 351.332 \pm 0.002$ s. Combined with previous X-ray and optical measurements of the spin period of the white dwarf, we find that its spin-up rate $\dot{P} = 0.00027 \pm 0.00005$ s yr⁻¹.

1. Introduction

GK Per is an extraordinary cataclysmic variable: the first nova of the last century (Nova Per 1901), it contains a magnetic white dwarf, a large truncated accretion disk, and an evolved secondary; it is surrounded by a nova shell visible in the radio, optical, and X-ray wavebands; and it suffers large-amplitude ($\Delta V \approx 3$), long ($\delta t \approx 60$ days), but infrequent ($\Delta t \approx 900$ day) dwarf nova outbursts (Simon 2002). On the strength of proposals submitted to the *XMM* and *Chandra* Guest Observer programs, we were granted target-of-opportunity *XMM* and *Chandra* (C. Mauche, PI) and *RXTE* (K. Mukai, PI) observations of GK Per during its 2002 March–April outburst with the goals of monitoring the X-ray light curve, pulse profile, and spectrum and obtaining the first grating spectra of GK Per to determine the nature of its complex X-ray spectrum (Sen & Osborne 1998; Ezuka & Ishida 1999). The number and dates of the observations were constrained by the small and decreasing angle between GK Per and the Sun, but we were fortunate to obtain *RXTE* monitoring observations every few days throughout the outburst, an *XMM* observation on March 9 during the early rise to maximum, and *Chandra* HETG observations on March 27 and April 9 during the late rise and peak of the outburst. The *XMM* MOS and *Chandra* HETG confirm that the Fe 6.4 keV fluorescent emission line is the strongest of the three Fe K-shell emission features, and the *XMM* RGS and *Chandra* HETG spectrometers reveal emission lines of H-like C; H- and He-like N, O, Ne, Mg, Si, and S; and possibly He-like Al (see Mukai et al. 2003 for a plot of the *Chandra* HETG spectrum). At least two sources of X-ray emission are implied by (1) the strong absorption of the continuum (which diminishes to zero for $\lambda \gtrsim 6$ Å), (2) the presence of emission lines down to $\lambda \approx 30$ Å, and (3) the fact that the short-wavelength ($\lambda = 1\text{--}6$ Å) flux is modulated at the spin period of the white dwarf, while the long-wavelength ($\lambda = 6\text{--}26$ Å) flux is steady.




Figure 1. Power spectrum of the *Chandra* HETG first-order 1–6 Å light curve of GK Per. The peaks at $\nu = 2.85$ and 5.68 mHz are identified with the fundamental and first harmonic of the white dwarf spin period $P = 351$.

2. Spin Period of the White Dwarf

To determine the spin period of the white dwarf of GK Per, we used data from the two 32 ks *Chandra* observations separated by 13.2 days. Specifically, we created a count rate light curve of the *Chandra* HETG first-order $\lambda = 1\text{--}6$ Å event data with $351.3/5=70$ s bins, subtracted the mean of the two observations, and padded the interval between the two observations with zeros. The power spectrum of the resulting light curve is shown in Figure 1, with peaks at $\nu = 5.68$, 2.85, 0.59, and 0.18 mHz, corresponding to periods $P = 176$, 351, 1700, and 5520 s; the 351 s period is the spin period of the white dwarf P , the 176 s period is its first harmonic $P/2$, and the 5520 s period is the quasi-periodic oscillation discussed by Morales-Rueda, Still, & Roche (1999). Based solely on the power spectrum, the spin period of the white dwarf can be constrained only to lie in the range $P = 349\text{--}353$ s, but by phasing the two *Chandra* HETG light curves, the number of cycles between the two observations is constrained to be $3250 \pm$ a few, corresponding to periods $P = 351.332$ s \pm multiples of 0.105 s. This ambiguity is resolved by the light curves of the two intervening *RXTE* PCA observations, which phase consistently with the two *Chandra* HETG light curves only for $P = 351.332 \pm 0.002$ s (see Fig. 2).

3. Spin-Up Rate of the White Dwarf

The pulse period of GK Per has been measured previously in X-rays by *Einstein* (Eracleous, Patterson, & Halpern 1991) and *EXOSAT* (Watson, King, & Osborne 1985; Norton, Watson, & King 1988) and in the optical by Patterson (1991), who provides an excellent discussion of the measurements and uncertainties of the various determinations of this quantity. Patterson finds that the 1980 *Einstein* data are consistent with pulse periods $P = 351.3423$ s \pm multiples of 0.002 s, the 1983–85 *EXOSAT* data are consistent with pulse periods $P = 351.3372$, 351.3409, or 351.3446 (± 0.0007) s, and the 1989–91 optical data




Figure 2. Phase-folded X-ray light curves of the first and second *Chandra* HETG and the intervening eighth and ninth *RXTE* PCA observations of GK Per. Data are shown by the histograms and the scaled mean phase-folded light curve is shown by the gray curves. Phase zero corresponds to HEJD 2452360.9113.




Figure 3. Spin period history of GK Per. The solid line is the spin-up rate $\dot{P} = 0.00027 \text{ s yr}^{-1}$ consistent with the *Einstein*, *EXOSAT*, optical, and *Chandra* and *RXTE* data, and the dotted line is the value $\dot{P} = 0.00084 \text{ s yr}^{-1}$ proposed by Patterson (1991).

are consistent with a pulse period $P = 351.3355 \pm 0.0004$ s; these data are plotted in Fig. 3. Assuming that the pulse period at the epoch of the *EXOSAT* observations was 351.3409 s, Patterson proposed a spin-up rate $\dot{P} = 0.0008$ s yr⁻¹ (the dotted line in Fig. 3) for the white dwarf of GK Per, but cautioned that the result should be regarded as tentative, as there is an uncertainty in the cycle count. As shown in Figure 3, our determination of the pulse period of GK Per in 2002 is not consistent with a linear extrapolation of the spin-up rate proposed by Patterson, but is consistent with a spin-up rate $\dot{P} = 0.00027 \pm 0.00005$ s yr⁻¹ (the solid line in Fig. 3).

4. Spin Ephemeris of the White Dwarf

Given the results from the previous sections, the spin ephemeris of the white dwarf of GK Per is fully determined by specifying the date of some feature in the light curve. Following Watson, King, & Osborne (1985), we define phase zero to occur at pulse maximum, which, guided by the power spectrum, we determined by fitting a function $a + b \sin[2\pi(\phi - \phi_1)] + c \sin[4\pi(\phi - \phi_2)]$ to the mean phase-folded light curve. By this method, we find that X-ray pulse maximum of GK Per follows the ephemeris $T_{\max} = \text{HEJD } 2452360.9113 (\pm 0.0001) + 0.00406634 (\pm 0.00000002) E - (3.5 \pm 0.6) \times 10^{-14} E^2$. This ephemeris includes only the second secure determination of the spin period of the white dwarf of GK Per and the first secure determination of its spin-up rate.

Acknowledgments. This work benefited from discussions with J. Patterson, K. Mukai, and A. Norton. We thank J. Swank and H. Tananbaum for the grants of Director's Discretionary Time which made the *RXTE* and *Chandra* observations possible. Support for this work was provided in part by NASA through *Chandra* Award Number DD2-3014X issued by the *Chandra* X-Ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-39073. This work was performed under the auspices of the U.S. Department of Energy by University of California Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

References

- Eracleous, M., Patterson, J., & Halpern, J. 1991, *ApJ*, 370, 330
 Ezuka, H., & Ishida, M. 1999, *ApJS*, 120, 277
 Morales-Rueda, L., Still, M. D., & Roche, P. 1999, *MNRAS*, 306, 753
 Mukai, K., et al. 2003, *ApJ*, in press (astro-ph/0301557)
 Norton, A. J., Watson, M. G., & King, A. R. 1988, *MNRAS*, 231, 783
 Patterson, J. 1991, *PASP*, 103, 1149
 Şen, G., & Osborne, J. P. 1998, in *Wild Stars in the Old West*, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 463
 Šimon, V. 2002, *A&A*, 382, 910
 Watson, M. G., King, A. R., & Osborne, J. 1985, *MNRAS*, 212, 917