



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Status of Continuum Edge Gyrokinetic Code Physics Development

X. Q. Xu, Z. Xiong, M. R. Dorr, J. A. Hittinger, G. D.
Kerbel, W. M. Nevins, B. I. Cohen, R. H. Cohen

June 1, 2005

10th IAEA Technical Meeting on "H-Mode Physics and
Transport Barriers"
St. Petersburg, Russia
September 28, 2005 through September 30, 2005

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Status of Continuum Edge Gyrokinetic Code Physics Development*

X. Q. Xu, Z. Xiong, M. R. Dorr, J. A. Hittinger, G. D. Kerbel,
W. M. Nevins, B. I. Cohen, and R. H. Cohen
Lawrence Livermore National Laboratory, CA 94550

We are developing an edge gyro-kinetic continuum simulation code to study the boundary plasma over a region extending from inside the H-mode pedestal across the separatrix to the divertor plates. A 4-D ($\psi, \theta, \epsilon, \mu$) version of this code is presently being implemented, en route to a full 5-D version. A set of gyrokinetic equations[1] are discretized on computational grid which incorporates X-point divertor geometry. The present implementation is a Method of Lines approach where the phase-space derivatives are discretized with finite differences and implicit backwards differencing formulas are used to advance the system in time. A fourth order upwinding algorithm is used for particle cross-field drifts, parallel streaming, and acceleration. Boundary conditions at conducting material surfaces are implemented on the plasma side of the sheath. The Poisson-like equation is solved using GMRES with multi-grid preconditioner from HYPRE. A nonlinear Fokker-Planck collision operator from STELLA[2] in $(v_{\parallel}, v_{\perp})$ has been streamlined and integrated into the gyro-kinetic package using the same implicit Newton-Krylov solver and interpolating F and $dF/dt|_{coll}$ to/from (ϵ, μ) space.

With our 4D code we compute the ion thermal flux, ion parallel velocity, self-consistent electric field, and geo-acoustic oscillations, which we compare with standard neoclassical theory for core plasma parameters; and we study the transition from collisional to collisionless end-loss. In the real X-point geometry, we find that the particles are trapped near outside midplane and in the X-point regions due to the magnetic configurations. The sizes of banana orbits are comparable to the pedestal width and/or the SOL width for energetic trapped particles. The effect of the real X-point geometry and edge plasma conditions on standard neoclassical theory will be evaluated, including a comparison of our 4D code with other kinetic neoclassical codes (such as NCLASS[3] and XGC[4]) and experiments.

References

- [1] T. S. Hahm, Phys. Plasmas , Vol. 3, 4658(1996).
- [2] K. Kupfer, R. W. Harvey, O. Sauter, et al., Phys. Plasmas **3**, 3644 (1996).
- [3] W. A. Houlberg, K. C. Shaing, S. P. Hirshman, M. C. Zarnstorff, Phys. Plasmas **4**, 3230(1997).
- [4] C. S.Chang, Seunghoe Ku, and H. Weitzner, Phys. Plasmas, Vol. 11, 2649(2004).

*This work was performed under the auspices of the USDoE by University of California Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.