



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

QED and electron collisions in the super strong fields of K-shell actinide ions

P. Beiersdorfer

February 1, 2006

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

03-ERD-004 Final Report

QED and electron collisions in the super strong fields of K-shell actinide ions

Principle Investigator:Peter Beiersdorfer
DivisionV
Phone.....3-3985
Email....beiersdorfer@llnl.gov
Mail StationL-260

Co-Investigators:.....Elmar Träbert (V Division)
.....Hui Chen (V Division)
.....Daniel Thorn (V Division)
.....Mau Chen (V Division)
.....Kennedy Reed (V Division)
.....Bret Beck (N Division)
.....John Becker (N Division)
.....Stephane Terracol (I Division)
.....Kwok-Tsang Cheng (A Division)
.....Richard Kelley (Goddard Space Flight Center)
.....Caroline Stahle (Goddard Space Flight Center)
.....Kevin Boyce (Goddard Space Flight Center)
.....Greg Brown (Goddard Space Flight Center)
.....Scott Porter (Goddard Space Flight Center)
.....Jonathan Sapirstein (Notre Dame University)

Objective: Atomic physics of high-Z, heavy ions is very different from that encountered in low-Z or medium-Z ions. The reason is the ultra strong nuclear field found only in the heaviest ions. The highest-Z atomic systems available to physical investigation, the actinides, therefore, offer rich new physics that cannot be studied any other way. This ranges from new dominating forces in electron-ion collisions to tests of fundamental theories. A measurement of the two-loop Lamb shift in uranium is by many considered to be the 'holy grail' of high-field QED tests of atomic systems. Such measurements have been attempted at heavy-ion accelerator facilities but have yet to succeed because of the difficulty to make measurements with the required accuracy. Also, electron collisions behave very differently in such tightly bound systems. The magnetic interaction between the ion and the incoming free electron (the so-called generalized Breit interaction) is essentially non-existent in collisions involving low and medium-Z ions. This interaction is therefore missing in essentially all electron collision codes. But in heavy, highly

charged ions like uranium, the generalized Breit interaction readily is the dominant force, changing electron collision cross sections by a factor of two. This has never been experimentally observed. In fact, no K-shell emission spectrum of any heavy high-Z ion higher than krypton ($Z=36$) has ever been recorded from a collisional source. By studying the heaviest actinides such fundamental science can be extended to regimes where the highest precision tests can be made.

Accomplishments: We successfully carried out our project. An overview is given below, and details can be found in references cited in the following.

- (1) We have resumed operation of the world's only source of stationary highly charged ions SuperEBIT [1]. This source is located at LLNL.
- (2) We have developed an injection system for introducing actinides into SuperEBIT. For this we implemented a 2J laser injection system that now routinely operates on SuperEBIT.
- (3) We have implemented a new high-energy microcalorimeter on SuperEBIT for x-ray and gamma ray measurements [2]. This calorimeter was built by our collaborators at NASA Goddard Space Flight Center and was delivered to us in October 2003. It has been used to measure the gamma rays produced by U-233 decaying into ^{229}Th , as well as from highly stripped U ions.
- (4) We have studied electron-ion collisions in the fully relativistic regime and tested magnetic sublevel calculations. [3,4]
- (5) We have made a series of measurements that progressively improved the accuracy of our QED measurements [4-9]. This series culminated in the best-ever QED measurement in highly charged uranium ions, which enabled us for the first time to isolate the two-loop QED contributions [10,11].

References:

- [1] Highly charged ion research at the Livermore electron beam ion traps
P. Beiersdorfer
Physica Scripta T120, 40-46 (2005)
UCRL-JC-152100
- [2] The Astro-E2 XRS/EBIT Microcalorimeter X-ray Spectrometer
F. S. Porter, G. V. Brown, K. R. Boyce, R. L. Kelley, C. A. Kilbourne, P. Beiersdorfer, H. Chen, S. Terracol, S. M. Kahn, A. E. Szymkowiak
Review of Scientific Instruments 75, 3772–3774 (2004)
UCRL-JRNL-203656
- [3] X-ray Line Polarization of the K-shell Resonance Line Emission of Heliumlike and Lithiumlike Sulfur at Relativistic Electron Impact Energies
D. L. Robbins, A. Ya. Faenov, T. A. Pikuz, H. Chen, P. Beiersdorfer, M. J. May, J. Dunn, K. J. Reed, and A. J. Smith
In *Proceedings of the 4th US-Japan Workshop on Plasma Polarization Spectroscopy, Kyoto, Japan, February 4-6, 2004*, T. Fujimoto and P. Beiersdorfer, editors, NIFS Proceedings Series No. NIFS-PROC-57 (National Institute for Fusion Studies, Toki, Japan 2004), p. 97–105.
UCRL-JRNL-202750
- [4] Measurement of the Polarization of the K-shell Resonance Line Emission of S^{13+} and S^{13+} at Relativistic Electron Beam Energies
D. L. Robbins, A. Ya. Faenov, T. A. Pikuz, H. Chen, P. Beiersdorfer, M. J. May, J. Dunn, K. J. Reed, and A. J. Smith
Physical Review A 70, 022715 (2004)
UCRL-JRNL-203478
- [5] Flat-field Grating Spectrometer for High-Resolution Soft X-ray and EUV Measurements on an Electron Beam Ion Trap
P. Beiersdorfer, E. Magee, E. Träbert, H. Chen, J. K. Lepson, M.-F. Gu, M. Schmidt
Review of Scientific Instruments 75, 3723-3726 (2004)
UCRL-JRNL-203237
- [6] Extreme ultraviolet spectra of highly charged Xe ions
E. Träbert, P. Beiersdorfer, J. K. Lepson, and H. Chen
Physical Review A 68, 042501 (2003)
UCRL-JC-153669

[7] Laboratory wavelengths of K-shell resonance lines of O V and O VI
M. Schmidt, P. Beiersdorfer, H. Chen, D.B. Thorn, E. Träbert, E. Behar
Astrophysical Journal 604, 562–564 (2004)

UCRL-JRNL-201118

[8] Wavelengths of the $4s_{1/2} - 4p_{2/3}$ resonance lines in Cu- and Zn-like heavy ions
E. Träbert, P. Beiersdorfer, H. Chen
Physical Review A 70, 032506 (2004)

UCRL-JRNL-2002764

[9] Soft-x-ray spectra of highly charged Os, Bi, Th, and U ions in an electron beam ion trap
E. Träbert, P. Beiersdorfer, K. B. Fournier, M. H. Chen
Canadian Journal of Physics 83, 829 (2005)

UCRL-JRNL-208760

[10] Testing high-Z QED with SuperEBIT: An estimate of the U^{91+} 1s two-loop Lamb shift
based on a measurement of the $2s_{1/2} - 2p_{1/2}$ transition in U^{89+}

P. Beiersdorfer

Radiation Physics and Chemistry

UCRL-JRNL-217517

[11] Measurement of the two-loop Lamb shift in lithiumlike U^{89+}

P. Beiersdorfer, H. Chen, D. B. Thorn, E. Träbert

Physical Review Letters 95, 233003 (2005)

UCRL-JRNL-211310

Acknowledgment: This work was performed under the auspices of the U.S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. The project 03-ERD-004 was funded by the Laboratory Directed Research and Development Program at LLNL.