



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

The XMM Cluster Survey: A Massive Galaxy Cluster at $z =$ 1.45

S. A. Stanford, A. K. Romer, K. Sabirli, M. Davidson, M. Hilton,
P. T. P. Viana, C. A. Collins, S. T. Kay, A. R. Liddle, R. G.
Mann, C. J. Miller, R. C. Nichol, M. J. West, C. J. Conselice, H.
Spinrad, D. Stern, K. Bundy

June 14, 2006

The Astrophysical Journal

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

The XMM Cluster Survey: A Massive Galaxy Cluster at $z = 1.45$

S.A. Stanford^{1,2}

A. Kathy Romer³

Kivanc Sabirli⁴

Michael Davidson⁵

Matt Hilton⁶

Pedro T.P. Viana^{7,8}

Chris A. Collins⁶

Scott T. Kay⁹

Andrew R. Liddle³

Robert G. Mann⁵

Christopher J. Miller¹⁰

Robert C. Nichol¹¹

Michael J. West^{12,13}

Christopher J. Conselice¹⁴

Hyron Spinrad¹⁵

Daniel Stern¹⁶

Kevin Bundy¹⁷

ABSTRACT

We report the discovery of XMMXCS J2215.9-1738, a massive galaxy cluster at $z = 1.45$, which was found in the *XMM Cluster Survey*. The cluster candidate was initially identified as an extended X-ray source in archival XMM data. Optical spectroscopy shows that 6 galaxies within a ~ 60 arcsec diameter region lie at $z = 1.45 \pm 0.01$. Model fits to the X-ray spectra of the extended emission yield $kT = 7.4_{-1.8}^{+2.7}$ keV (90% confidence); if there is an undetected central X-ray point source then $kT = 6.5_{-1.8}^{+2.6}$ keV. The bolometric X-ray luminosity is $L_x = 4.4_{-0.6}^{+0.8} \times 10^{44}$ ergs s $^{-1}$ over a 2 Mpc radial region. The measured T_x , which is the highest for any known cluster at $z > 1$, suggests that this cluster is relatively

¹University of California, Davis, CA 95616; adam@igpp.ucllnl.org

²Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, Livermore, CA 94551

³Astronomy Centre, University of Sussex, Falmer, Brighton, BN1 9QJ, U.K.

⁴Department of Physics, Carnegie Mellon University, 5000 Forbes Ave., Pittsburgh, PA, 15217

⁵Institute of Astronomy, University of Edinburgh, Blackford Hill, Edinburgh, EH9 9HJ, U.K.

⁶Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkinhead CH41 1LD, U.K.

⁷Departamento de Matemática Aplicada da Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre, 687, 4169-007, Portugal

⁸Centro de Astrofísica da Universidade do Porto, Rua das Estrelas, 4150-762, Porto, Portugal

⁹Astrophysics, Keble Road, Oxford, OX1 3RH, U.K.

¹⁰Cerro-Tololo Inter-American Observatory, National Optical Astronomy Observatory, 950 North Cherry, Tucson, AZ, 85719

¹¹Institute of Cosmology and Gravitation, University of Portsmouth, Portsmouth, PO1 2EG, U.K.

¹²Department of Physics and Astronomy, University of Hawaii, 200 W. Kawili Street, Hilo, Hawaii 96720, U. S. A.

¹³Gemini Observatory, Casilla 603, La Serena, Chile

¹⁴University of Nottingham, University Park, Nottingham, NG9 2RD, UK

¹⁵Astronomy Department, University of California, Berkeley, CA 94720

¹⁶Jet Propulsion Laboratory, California Institute of Technology, MS 169-506, 4800 Oak Grove Road., Pasadena, CA, 91109

¹⁷California Institute of Technology, MS 105-24, 1200 California Blvd., Pasadena, CA, 91125

massive for such a high redshift. The redshift of XMMXCS J2215.9-1738 is the highest currently known for a spectroscopically-confirmed cluster of galaxies.

Subject headings: galaxies: clusters — galaxies: evolution — galaxies: formation

1. Introduction

High redshift galaxy clusters provide important laboratories for the study of structure formation and galaxy evolution. They also can be used to constrain cosmological parameters independent of the Cosmic Microwave Background and supernova methods. The number of known clusters at $z > 1$ is currently small ($\simeq 10$ with X-ray confirmation: e.g. Stanford et al. 2001, 2002; Rosati et al. 2004; Mullis et al. 2005), but growing rapidly thanks to new surveys being carried out in the X-ray (e.g. Romer et al. 2001; Mullis et al. 2005; Bremer et al. 2006; Barkhouse et al. 2006), optical (e.g. Gladders & Yee 2005), and infrared (e.g. Stanford et al. 2005). In the near future, Sunyaev-Zel'dovich surveys should also provide large numbers of high redshift clusters (Carlstrom et al. 2002). We announce here the discovery of a high redshift cluster in the XMM Cluster Survey (XCS: Romer et al. 2001) and the measurement of its X-ray temperature. At $z = 1.45$, XMMXCS J2215.9-1738 is the most distant cluster with spectroscopic redshift confirmation to date. Unless otherwise noted, we assume $\Omega_m = 0.27$, $\Lambda = 0.73$, and $H_o = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. Observations

2.1. X-ray

XMM-Newton is the most sensitive X-ray spectral-imaging telescope deployed to date. The XCS identifies cluster candidates in the *XMM* data archive using the signature of X-ray extent. With a projected total area of 500 deg^2 , the XCS is expected to catalog several thousand clusters out to a redshift of $z \simeq 2$ (Romer et al. 2001). To date, 1,847 *XMM* observations have been processed using an automated pipeline, resulting in a survey area of 168 deg^2 at $b > 20^\circ$. This is the net area available for cluster searching and excludes the Magellanic Clouds, artifacts due to out of time events, and X-ray sources that extend over a large fraction of the field of view, such as supernova remnants, Galactic clusters, or low redshift galaxy clusters. The initial catalog contains 1764 cluster candidates (see Davidson et al. in preparation and Sabilri et al. in preparation for details). Here we focus on 5 moderately deep observations of varying length in the direction of the $z = 2.217$ quasar

LBQS 2212-1759, which were obtained in 2000 and 2001. An extended source was detected at the 4σ or greater level by the XCS pipeline at the same location, $9'$ from the center of the field of view, in 3 of the 5 observations. The combined *XMM* exposure time, after correction for solar flares etc., in the direction of the cluster candidate is 237 ks (MOS1). The typical point source positional uncertainty in the XCS is 3 arcsec at 9 arcmin off-axis (Davidson et al. in preparation) and we have verified this by checking the positions of three quasars in the summed X-ray data near the cluster candidate.

2.2. Optical and IR Imaging

An *I*-band image of the area around XMMXCS J2215.9-1738 was found in the ESO Imaging Survey (Dietrich et al. 2005). This 9000 s image reaches a limiting magnitude of 24 (Vega), and allows even $z > 1$ early-type galaxies to be detected. On the basis of visual inspection of the *I*-band image, XMMXCS J2215.9-1738 was selected as a possible $z > 1$ galaxy cluster, worthy of spectroscopic followup.

Subsequent to the first slit mask spectroscopy (see below), K_s -band imaging was carried out on the 200 inch telescope at Palomar on 2005 October 21. A total of 3240 s of exposure time was obtained in clear conditions, and the images were reduced following standard infrared procedures. The reduced image was calibrated onto the Vega system by comparison of instrumental magnitudes for stars with the same objects in the 2MASS database. The image is shown in Figure 1, along with contours of the X-ray emission. The X-ray emission from the cluster candidate is clearly extended compared to the three nearby point sources. None of the X-ray sources in Figure 1 (the cluster, the three point sources, or the southern extension) has been cataloged previously, based on an examination of the NED and SIMBAD databases.

The *I* and K_s band images were photometered using SExtractor. Objects were detected on the K_s image and then aperture magnitudes of diameter 2.5 arcsec were measured on both images using the same SExtractor catalog. The resulting color-magnitude diagram shown in Figure 2 was used for target selection for the second mask in the spectroscopic followup program.

2.3. Keck Spectroscopy

The first spectroscopic followup was carried out using the Deep Imaging Multi-Object Spectrograph (DEIMOS: Faber et al. 2003) at the Keck II telescope on UT 2005 September

1. Targets were selected from the I -band image to have $23 < I < 24$ and be located in the central few arcmin of the area around the X-ray candidate cluster. The slits had widths of 1.0 arcsec and minimum lengths of 5 arcsec. We used the 600ZD grating, which is blazed at 7500 Å, to cover a nominal wavelength range of 5000 to 10000 Å. The dispersion of ~ 0.65 Å pixel $^{-1}$ resulted in a spectral resolution of 3.8 Å. Of particular note is that this spectral resolution is adequate for being able to detect both lines in the [O II] λ , 3727 doublet. We obtained 5×1800 s exposures with this setup in photometric conditions with 1.1 arcsec seeing. The observations were carried out with the slitlets aligned close to the parallactic angle.

A second slit mask was observed, again using DEIMOS at the Keck II telescope, on UT 2005 November 5. For this mask, we were able to use the color information seen in Figure 2 to select targets likely to be in the cluster by choosing objects in the area of the color-magnitude diagram where the red sequence is expected for a $z = 1.45$ cluster. These observations were conducted in poor conditions, with ~ 1.5 arcsec seeing. A total of 5400 s of exposure time was obtained with the second mask.

The slitmask data for the two masks were separately reduced using the DEEP pipeline. A relative flux calibration was obtained from longslit observations of the standard stars G191B2B and Wolf 1346. One-dimensional spectra were extracted from the sum of the reduced data for each slitlet.

3. Results

3.1. X-ray Analysis

To examine the X-ray emission from the intracluster medium, we need to account for contamination from the nearby point sources. At the same time it behooves us to allow for a possible central point source, such as a galaxy with an active galactic nucleus, that would affect our attempts to extract information on e.g. the temperature of the ICM. Using SHERPA, we carried out 2-dimensional fitting of the detected nearby point sources simultaneously with fitting of the extended emission from the cluster candidate. In addition to the obvious point sources to the east of the cluster, we have included a variable strength point source fixed at the center of the extended emission. The profile of the X-ray emission is shown in Figure 3. The best fitting β model combined with the point sources is shown, along with a PSF which indicates that extended emission is present. The brightest point sources are obvious as bumps in the profile at radii of ~ 37 , 60, and 75 arcsec. We find that a central point source contributes at most 10% of the counts within the central $r = 30$ arcsec region.

The best fitting β model for the extended component has a core radius of $r_c = 73_{-16}^{+18}$ kpc and $\beta = 0.53_{-0.03}^{+0.04}$.

Next we used the results of this two-dimensional fitting procedure to accurately determine the spectrum of the extended emission after correction for contamination by the point sources. The appropriate counts due to the point sources were included (as fixed models normalized to the results of the spatial analysis) while carrying out the fitting of the extended source spectrum as described below. We have extracted a spectrum for the cluster candidate using photons detected within a $30''$ radius of $\alpha = 22^h15^m58^s.5$, $\delta = 17^\circ38'2''.5$. We extracted MOS1, MOS2 and pn spectra from each XMM observation with an exposure time > 10 ks made towards the cluster. We generated a combined spectrum for each camera and these are presented in the left panel of Figure 4. Spectral fitting was carried out using XSPEC v11.3.1s (Arnaud 1996) using the MEKAL models (Mewe et al. 1986) modified with an interstellar absorption (McCammon & Sanders 1990) appropriate for the Galactic column density at that location. A local background correction was made using a spectrum extracted from a $30''$ radius region at the same off-axis angle ($9'$) as the cluster; this avoids having to carry out an energy dependent vignetting correction of the background. To find the best fit spectrum, we limited the fit to the energy range $0.5 - 7$ keV and grouped the data so that there were a minimum of 20 counts per bin. The number of background corrected counts in that energy range (summed across all three cameras) was 1100 in the $30''$ measurement aperture, after correction for the point sources. We fixed the hydrogen column density to the Dickey & Lockman (1990) value, the metal abundance to $Z = 0.3$ times the Solar value and, initially, the redshift to the Keck spectroscopic value, $z = 1.45$. After accounting for all point sources including the possible central source, the best fit temperature of the extended emission was found to be $T_x = 6.5_{-1.8}^{+2.6}$ keV (90% confidence limits). If there is no central point source, then the best fit temperature of the ICM would be $T_x = 7.4_{-1.8}^{+2.7}$ keV. The best fit model, and its residuals, are shown in Figure 4. Using this spectrum, we have determined the cluster flux to be $1.22_{-0.17}^{+0.21} \times 10^{-14}$ erg s $^{-1}$ cm $^{-2}$ [0.5 -2.0 keV], extrapolated to a region 2 Mpc in radius using the best fit β model. For the same 2 Mpc aperture, the luminosity of the ICM is $1.15_{-0.16}^{+0.20} \times 10^{44}$ erg s $^{-1}$ in the [0.5 - 2.0] keV band, and $4.39_{-0.61}^{+0.76} \times 10^{44}$ erg s $^{-1}$ in a bolometric band [0.05 - 100 keV].

3.2. Optical Spectroscopy

Visual inspection of the reduced spectra from both DEIMOS masks yielded redshifts for objects. The 6 galaxies listed in Table 1 lie within 30 arcsec of the X-ray centroid and have concurrent redshifts $1.445 < z_{spec} < 1.455$. In 3 of the 6 spectra, the [O II] λ 3727

doublet is detected and clearly split. The other three objects have spectra which show features from the redshifted 4000Å and 2900Å breaks. The spectrum of one of the cluster members, ID 14339, is shown in Figure 5. In addition to these 6 galaxies we consider to be members of the cluster, three more galaxies were spectroscopically identified with redshifts at $1.44 < z < 1.46$. However, they lie approximately 5 – 7 arcmin from the X-ray centroid, so are not (yet) clearly members of the cluster.

3.3. Optical and Infrared Photometry

The color–magnitude diagram for all objects in a 3′ area at the cluster is shown in Figure 2. The objects within 30 arcsec of the X-ray centroid are circled in red. The 6 galaxies with concurrent spectroscopic redshifts $1.445 < z < 1.455$ are shown by the solid red points. The scatter seen in the colors of the member galaxies is dominated by the photometric uncertainties in the photometry so is not indicative of the intrinsic scatter in the red sequence in this cluster. Also plotted in Figure 2 are two estimates of the expected color of L* early–type galaxies. Using a simple passive evolution model calculated from the GISSEL of Bruzual & Charlot (2003) and our assumed cosmology, we calculated the expected colors for a single 100 Myr burst stellar population of solar metallicity stars formed at $z_f = 3.0$ and $z_f = 2.0$, with slopes assumed to be the same as those in Coma for a similar rest frame color. As can be seen in Figure 2, the colors of the member galaxies are consistent with those expected for early-type galaxies whose stars formed at $2 < z_f < 3.0$.

4. Discussion

Analyses of our combined X-ray imaging spectroscopy, optical spectroscopy, and optical-IR imaging provide strong evidence that XMMXCS J2215.9-1738 is a massive galaxy cluster at $z = 1.45$. At least 10 galaxy clusters with $z > 1.4$ and $T_x > 6$ keV are expected to be present in our current XCS area of 168.2 deg², if the universe is assumed to be flat with $\Omega_m = 0.238$ and $\sigma_8 = 0.744$ (WMAP - Spergel et al. 2006). This estimate requires the relation between cluster mass and X-ray temperature to be normalized so as to reproduce the local abundance of X-ray clusters (HIFLUGCS - Reiprich & Böhringer 2002). However, it should be noted that these assumptions require that a local 5 keV cluster would need to have a mass, within an overdensity of 500 with respect to the critical density, of around $2 \times 10^{14} h^{-1} M_\odot$ (Pierpaoli et al. 2003; Viana et al. 2003), about 2/3 smaller than is implied by the most recent observational data (Arnaud, Pointecouteau & Pratt 2005; Vikhlinin et al. 2006).

With the combination of the serendipitous searches for clusters in the *XMM* archival data (Romer et al. 2001; Mullis et al. 2005), and the on-going wide-area surveys such as UKIDSS (Lawrence et al. in preparation) and the RCS (Gladders & Yee 2005) the time finally is ripe for the identification of large samples of $z > 1$ clusters. The construction of such samples will pave the way towards a better understanding of the origin of galaxy clusters and their constituent galaxy populations.

This work is based on data obtained by XMM-Newton, an ESA science mission funded by contributions from ESA member states and from NASA. We acknowledge financial support from; the NASA-LTSA program, the RAS Hosie Bequest, Liverpool John Moores University, the Institute for Astronomy at the University of Edinburgh, the XMM & Chandra guest observer programmes, Carnegie Mellon University, the NSF and PPARC. This research made use of the NASA/IPAC Extragalactic Database, the SIMBAD facility at CD-S, the NASA/GSFC supported XSPEC software, and the ESO Imaging Survey. The W. M. Keck Observatory is a scientific partnership between the University of California and the California Institute of Technology, made possible by a generous gift of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community; we are fortunate to have the opportunity to conduct observations from this mountain. The analysis pipeline used to reduce the DEIMOS data was developed at UC-Berkeley with support from NSF grant AST-0071048. We thank the referee for helping to improve the final version of this paper. The work by SAS at LLNL was performed under the auspices of the U.S. Department of Energy under Contract No. W-7405-ENG-48. The work of D.S. was carried out at the JPL, California Institute of Technology, under contract with NASA.

This work was performed in part under the auspices of the U. S. Department of Energy by University of California, Lawrence Livermore National Laboratory under contract W-7405-Eng-48.

REFERENCES

- Arnaud, K. in *Astronomical Data Analysis Software and Systems V*, 1996, A.S.P. Conf. Ser., 101, G. Jacoby & J. Barnes, eds., 17
- Arnaud, M., Pointecouteau, E., Pratt, G. W., 2005, *Å*, 441, 893
- Barkhouse, W. et al. 2006, *ApJ*, in press
- Bremer, W. et al. 2006, *MNRAS*, in press
- Bruzual, G. & Charlot, S. 2003, *MNRAS*, 344, 1000
- Carlstrom, J. E., Holder, G. P., & Reese, E. D. 2002, *ARA&A*, 40, 643
- Dickey, J.M. & Lockman, F.J., 1990, *ARA&A*, 28, 215
- Dietrich, J.P. et al. 2005, *astro-ph/0510223*
- Faber, S. et al. 2003, *SPIE*, 4841, 1657
- Gladders, M. & Yee, H. 2005, *ApJS*, 157, 1
- McCammon, D. & Sanders, D.T. 1990, *ARA&A*, 28, 657
- Mewe, R., Lemen, J.R. & van der Oord, G.H.J., 1986, *A & AS*, 62, 197
- Mullis, C. et al. 2005, *ApJ*, 623, L85
- Pierpaoli, E., Borgani, S., Scott, D., & White, M., 2003, *MNRAS*, 342, 163
- Reiprich, T. H. & Böhringer, H., 2002, *ApJ*, 567, 716
- Romer, K., Viana, P.T.P, Liddle, A.R., & Mann, R.G. 2001, *ApJ*, 547, 594
- Rosati, P. et al. 2004, *AJ*, 127, 230
- Spergel, D.N. et al., 2006, submitted to *ApJ*, *astro-ph/0603449*
- Stanford, S.A., et al. 2001, *ApJ*, 552, L504
- Stanford, S.A., Holden, B., Rosati, P., Eisenhardt, P., Stern, D., Squires, G., & Spinrad, H. 2002, *AJ*, 123, 619
- Stanford, S.A. et al. 2005, *ApJ*, 634, L129

Viana, P. T. P., Kay, S. T., Liddle, A. R., Muanwong, O., Thomas, P. A., 2003, MNRAS, 346, 319

Vikhlinin, A., Kravtsov, A., Forman, W., Jones, C., Markevitch, M., Murray, S. S., Van Speybroeck, L., 2006, ApJ, 640, 661

Table 1. Summary of Spectroscopic Cluster Members

ID	R.A. ^a	Dec. ^a	K_s	$I - K_s$	spec-z
14651	22:15:58.364	-17:37:37.5	>19.9	<4.0	1.4526
14478	22:15:57.233	-17:37:53.1	19.58	4.20	1.4537
14378	22:15:58.879	-17:37:59.6	19.49	4.00	1.451
14289	22:15:58.488	-17:38:10.4	19.04	4.49	1.453
14339	22:15:59.057	-17:38:02.0	19.18	4.16	1.4467
14389	22:15:58.479	-17:37:58.6	19.07	4.17	1.452

^aCoordinates are J2000.




Fig. 1.— K_s -band image of a 3 arcmin field centered on the cluster with an X-ray overlay of contours in blue. The 6 spectroscopically confirmed members are circled in red (ID=14651 is not detected in the K_s image). The green circle of diameter 60 arcsec shows the location of the region used for the X-ray analysis reported in section 3.1.




Fig. 2.— An $I - K_s$ vs K_s color-magnitude diagram for XMMXCS J2215.9-1738; aperture magnitudes are used for the colors and K_s magnitudes. The black points are all the objects within a 3 arcmin box around the position of the X-ray source, and the objects circled in red lie within 30 arcsec of the center of the X-ray source. The solid red points are those with concurrent spectroscopic redshifts at $z = 1.45$. The dashed (dotted) line represents the expected color-magnitude relation for galaxies as predicted by a Bruzual & Charlot model in which all the stars formed within a 0.1 Gyr burst starting at $z_f = 3$ ($z_f = 2$) in the standard cosmology. The solid lines represent the 5σ limits to the K_s and $I - K_s$ photometry.




Fig. 3.— The spatial profile of the X-ray emission. The points are the data. The best fitting model, combining both the extended component and the point sources (which are obvious as bumps at large radii), is shown by the solid line. A PSF is shown by the dashed line. The background is shown by the horizontal dotted line.




Fig. 4.— X-ray spectrum of XMMXCS J2215.9-1738. Counts from EMOS1, EMOS2, and EPN are plotted with black, red, and green respectively. The spectra are after accounting for the point sources, including a possible central source. The best fit to these spectra is an absorbed MEKAL model with properties specified in the text; the folded model is overplotted (only T_x and the normalisation are left as free parameters).




Fig. 5.— An optical spectrum of cluster member 14339 with $z = 1.45$. The identified spectral features are marked. The spectrum has been smoothed by a 6.5\AA boxcar.