



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Analysis of Radiocarbon, Stable Isotopes and DNA in Teeth to Facilitate Identification of Unknown Decedents

K. Alkass, H. Saitoh, B. A. Buchholz, G.
Holmlund, D. R. Senn, K. L. Spalding, H. Druid

August 21, 2012

PLOS ONE

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Analysis of radiocarbon, stable isotopes and DNA in teeth to facilitate identification of unknown decedents

Alkass K¹, Saitoh H², Buchholz BA³, Holmlund G⁴, Senn DR⁵, Spalding KL⁶, Druid H¹

¹Division of Forensic Medicine, Department of Oncology-Pathology, Karolinska Institutet, Retzius v. 3, SE-171 77 Stockholm, Sweden,

²Department of Legal Medicine, Graduate School of Medicine, Chiba University, 1-8-1 Inohana, Chou-ku, Chiba, 260-8670, Japan,

³Center for Accelerator Mass Spectrometry, Lawrence Livermore National Laboratory, 7000 East Avenue, L-397, Livermore, CA 94551, USA,

⁴Department of Forensic Genetics and Forensic Toxicology, National Board of Forensic Medicine, Linköping, Sweden, ⁵Center for Education and Research in Forensics, The University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229, USA,

⁶Department of Cell and Molecular Biology, Karolinska Institutet, Berzelius v. 3, SE-171 77 Stockholm, Sweden.

Corresponding author: Kanar Alkass; Division of Forensic Medicine, Department of Oncology-Pathology, Karolinska Institutet, Retzius v. 3, SE-171 77 Stockholm; +46 8 52487817. E-mail: kanar.alkass@ki.se

Classification: BIOLOGICAL SCIENCES (medical sciences)

Keywords: carbon-14, stable isotopes, teeth, age determination, forensic

Abstract

The characterization of unknown dead bodies or suspected human remains is a frequent and important task for forensic investigators. However, any identification method requires clues to the person's identity to allow for comparisons. If such clues are lacking, information about the year of birth, sex and geographic origin of the victim, is particularly helpful to aid in the identification casework and limit the search for possible matches. We here present results of stable isotope analysis of ^{13}C and ^{18}O , and bomb-pulse ^{14}C analyses that can help in the casework. The ^{14}C analysis of enamel provided information of the year of birth with an average absolute error of 1.8 ± 1.3 years. We also found that analysis of enamel and root from the same tooth can be used to determine if the ^{14}C values match the rising or falling part of the bomb-curve. Enamel laydown times can be used to estimate the date of birth of individuals, but here we show that this detour is unnecessary when using a large set of crude ^{14}C data of tooth enamel as a reference. The levels of ^{13}C in tooth enamel were higher in North America than in teeth from Europe and Asia, and Mexican teeth showed even higher levels than those from USA. DNA analysis was performed on 28 of the teeth, and provided individual-specific profiles in most cases and sex determination in all cases. In conclusion, these analyses can dramatically limit the number of possible matches and hence facilitate the identification work.

Introduction

The identification of unknown bodies is crucial for ethical, medico-legal and civil reasons. The police and forensic investigators worldwide spend considerable time every year attempting to solve such cases. Typically, unknown dead bodies or human remains are eventually identified by fingerprint comparisons, comparison of ante- and postmortem dental or medical radiographs, or by DNA analysis of tissue from the dead body and from possible relatives. However, in cases where there are no clues as to the identity, a characterization of the dead body can limit and focus the search for possible matches and help to exclude impossible or unlikely alternatives. The Doe Network Database (<http://www.doenetwork.org>) is extensively used in North America and includes both reported missing persons and unknown dead bodies. This database contains information on about 3,800 unidentified dead bodies, the majority from the United States. A careful review of this database (Table 1) showed that the estimated

age for adult individuals displayed an average uncertainty range of 15 ± 12 years. Such a wide range is not very helpful in the efforts to limit the search for possible matches. Furthermore, the sex was unknown or uncertain in some of the cases, and the geographic origin of the person rarely known, apart from the information about the place where the victim was found. Hence, a more detailed characterization of the subjects in terms of age, sex and geographical origin should be expected to improve the success rate in dead victim identification work.

The most common procedure to estimate the age of an unknown body is to perform an anthropological examination, which regarding skeletonized bodies focuses on age-related changes in teeth and bones. However, the anthropological methods are not very precise and typically an error of ± 10 years is added to the estimated age (1). Additional methods have therefore been developed. Today, the most precise method to determine the age at death is aspartic acid racemization analysis, which is based on the observation that remaining aspartic acid in the tooth trapped during its formation will be converted at a very slow rate from the L-form to the D-form, both of which can be detected and quantified by gas chromatographic methods (2). These methods applied on enamel, dentin or cementum can provide an estimate of the age of a person with a fairly good precision; using dentin the error may be ± 5 years or less (3-5). All the methods used to determine age are based on age-dependent alterations in human tissues, and hence, they will all give an estimate of the person's age at death. In contrast, the recently developed method to analyze bomb-pulse ^{14}C in enamel indicates the year of birth regardless of when, or at what age, the person died (6). This method should not be confused with the Libby method (7) to date archeological material, which is based on the radioactive decay of ^{14}C in biological material. Instead, the bomb-pulse ^{14}C method takes advantage of the above ground nuclear test bomb detonations 1955-1963, which caused a substantial increase in global ^{14}C levels (8-12). Repeated measurements of ^{14}C in the atmosphere and in biological products of known age, has over time resulted in reference values to which analytical results can be compared to offer an estimate of the age; see <http://calib.qub.ac.uk/CALIBomb>, (10).

Similarly, an important factor for limiting the search for possible matches is the sex of the individual. Unknown human bodies that are fairly well preserved rarely pose any problem, but if the forensic case involves a mutilated or skeletonized body, anthropological examination of bones that show sex dimorphism is often performed.

DNA profiling of bones and teeth is possible, and can include markers for sex. However, if a fire victim is charred, extraction and amplification of DNA from soft tissues and bones can be difficult. Teeth represent the most resistant tissue in the body and DNA analysis of teeth may be an alternative. Such analysis, including markers for sex, has successfully been performed even on incinerated bodies (13).

To limit the search for possible matches, the geographic origin of the deceased may also provide clues to the identity. In birds, analysis of stable isotopes in feathers formed during the winter has been used to determine wintering areas (14). In a recent study, hair samples from subjects living on different continents displayed geographically specific isotopic signatures (15). In a previous study, we have shown that such geographic differences regarding ^{13}C also can be detected in teeth (16). This isotope is incorporated into the tissues of animals, including humans, in relation to the content in food. Since the diet by tradition varies geographically, and is based on different primary products derived from plants with different ^{13}C content, the amounts in tissues will vary accordingly (17). The geographic differences can be explained as follows. Some types of plants can discriminate between ^{12}C and ^{13}C , resulting in differences in the levels of ^{13}C between different types of plants. C4 plants (which include corn and sugar cane) contain higher amounts of ^{13}C than C3 plants (which include potato, wheat and sugar beet) since these strive to maximize their CO_2 assimilation. In general, C4 plants tend to grow in hotter or drier climates than C3 plants. This in turn means that animals, including humans, having a diet based mainly of C4 plants and/or on animals that eat C4 plants, will incorporate more ^{13}C than those mainly living off C3 plant based diets.

^{18}O , is another stable isotope that shows geographic variation. The incorporation of ^{18}O in animal tissues is correlated to the levels in drinking water and these levels vary with latitude because of differences in the evaporation and condensation propensity between ^{16}O and ^{18}O (18). In the United States, reports indicate that the highest tap water ^{18}O concentrations are in the south and southeast and the lowest in the northwest (19). These data are based on tap water samples collected from 2002-2003. Due to differences in precipitation and variable ground water supplies to specific areas, these levels may differ somewhat from year to year. Still, ^{18}O levels are not expected to change dramatically over time and hence levels in teeth, bones and hair should mirror the levels in tap water fairly well.

In this study we investigated whether analysis of bomb-pulse derived ^{14}C levels in teeth from subjects in North America shows similar precision as in previous studies on teeth from other continents. In addition, we analyzed ^{14}C in both enamel and roots of the same teeth to find out if this measure could help to discriminate values that relate to incorporation during the rising or falling part of the bomb-curve. We also wanted to explore possible geographical variation in ^{13}C and ^{18}O by collecting teeth from Mexico, Canada and different parts of the United States. From the same teeth, we also performed DNA analysis of the amelogenin gene to determine the sex of the subjects. Finally, the average enamel ^{14}C incorporation time in different types of teeth from this study were combined with the results from previous studies (5, 6, 16) in order to provide a reference guide for determining the date of birth of persons.

Results

Date of birth (DOB) estimation using tooth enamel. For all 66 teeth, sufficient amounts of enamel were obtained to allow for ^{14}C determination by accelerator mass spectrometry (AMS). Table 2 displays the overall results and Table S1 provides more detailed information. Nine of the teeth had a calculated enamel formation time before 1955 and all showed pre-bomb ^{14}C values. For teeth laid down after 1955, a high correlation was found between ^{14}C levels in enamel and the actual formation time of the enamel, with an average absolute error of 1.8 ± 1.3 years, $R^2 = .9935$ ($n=57$). In 14 cases, two or more teeth were analyzed from the same individual. On average the mean difference in estimates between teeth from the same individuals was 1.4 (median 1.2) years. In Case 30, 14 teeth had been extracted. The ^{14}C values of each of these teeth predicted the true DOB very well and the difference in prediction of DOB between these teeth was small (Table 2). In nine of the twelve cases, where the enamel was formed after the onset of the bomb pulse, enamel ^{14}C values matched the expected order of formation, i.e. teeth with later formation time showed more recent ^{14}C levels. In Cases 9 and 19 both teeth from each individual showed very similar ^{14}C values since both tooth types had similar enamel formation times (Table 2). In Case 30, high ^{14}C values were seen in all teeth, which was in line with the fact that their actual formation dates were during the peak of the bomb curve.

^{14}C incorporation in tooth roots. We also analyzed ^{14}C levels in the roots from 28

teeth collected from 26 individuals (Table 3). The ^{14}C levels were consistently higher in the roots than in the enamel if the enamel was formed during the rising part of the curve, and lower in the roots if the enamel was formed during the falling part of the curve. One exception was Case 16, where the enamel was formed shortly before the peak of the bomb-spike and most likely the lower level of ^{14}C in the root is explained by lower levels incorporated during the falling part of the curve. In all pre-bomb cases, i.e. when the enamel was completely formed before the bomb pulse, post-bomb values were found in their corresponding roots, indicating that a turnover in some component of the root continues even into old age, as evidenced by the elevated ^{14}C concentration in Case 3, a man born in 1918. This finding confirms previous observations of a continuous formation of secondary dentin and cementum throughout life (20, 21).

Reference levels of ^{14}C to determine date of birth. The principle for DOB determination of persons using ^{14}C has been based on subtracting the enamel formation time for the particular type of tooth based on radiographic mapping of tooth development. However, in Table S2, we have compiled the enamel DOB as estimated by ^{14}C levels in this and previous studies (5, 6, 16) and compared the result for each type of tooth with the actual DOB of the person. We show that the error in birth dating can be reduced by subtracting the average enamel ^{14}C incorporation time instead.

^{13}C in teeth can tell geographic origin. The ^{13}C levels in the tooth enamel from 35 teeth (43 individuals) were analyzed with isotopic ratio mass spectrometry whenever the AMS processing allowed for a split (Table 4). The teeth extracted from subjects raised in North America showed higher enamel ^{13}C values compared to levels previously seen in teeth collected from other continents, including teeth from South America (16). The levels were somewhat lower in teeth from California, British Columbia and Connecticut, compared to teeth from Texas. Enamel from teeth from Mexico showed very high levels, even higher than levels in enamel from teeth from the United States and Canada, without overlap. ^{13}C levels in 31 tooth roots from 27 individuals were also analyzed; 19 of the roots were from the same teeth in which ^{13}C enamel levels also were measured (Table 4). The ^{13}C levels in the roots were generally lower than in the enamel. Furthermore, the ^{13}C levels in tooth roots were

higher in teeth from Mexico than in roots from the United States and Canada, but showed a higher variation, making it more difficult to separate Mexican subjects from persons raised in United States and Canada using roots only.

Tap water provides an additional signature. In Table 5, the ^{18}O levels in all roots analyzed are displayed. ^{18}O levels were in general lower in tooth roots from the northwestern region of North America (Washington and British Columbia) compared to those in roots collected from Texas ($p < 0.01$). One subject from Alabama and one subject from Texas were outliers, but otherwise the levels of ^{18}O paralleled the drink water levels fairly well (Fig. 1). Mexican teeth showed slightly lower values than teeth from Texas and were similar to the average for all teeth from United States and Canada. From Table 5, it can be appreciated that there are regional differences in the levels of ^{18}O , but at a lower magnitude as compared to the difference in tap water levels. Since we did not collect information about the exact place where the subjects were raised, interstate differences may explain the modest differences between states. Further, the levels of ^{18}O are also dependent of the diet, since a large part of the oxygen incorporated in tissues derive from food that may be produced at remote locations.

Sex determination. A small part of the root of 29 teeth from 25 individuals was used for DNA analysis. A full profile was obtained for 22 of the teeth and in all cases the sex could be determined using the amelogenin gene as a marker (22), see Fig. 2. In four cases, two teeth were analyzed and the results were identical for the different teeth from the same individual. Further, in select cases, the results of the amelogenin gene analysis was confirmed by analysis of Y chromosome specific STR markers using Powerplex® Y System (Promega Corp., Madison, WI).

Discussion

We report that ^{14}C analysis of enamel from teeth collected from North American subjects predicts the date of birth of the individual with an average absolute error of 1.8 ± 1.3 years. This precision is similar to that of previous studies on teeth obtained from other countries around the world (5, 6, 16, 23, 24). Given that a large number of above ground detonations were performed in the Nevada desert, a concern regarding the reliability of this method could be raised about subjects born and raised during the

cold war in nearby regions, such as California, Texas and northern Mexico. However, from Table 2 and Table S1, it can be appreciated that the precision of birth dating such individuals was not different than that of others in this study. The enamel formation times differ between different types of teeth, but also show a variation among the same type of tooth, with third molar teeth showing the highest variation (25, 26). Despite this fact, we found that analysis of two or more teeth, including third molars, from the same individual provided similar estimates of date of birth of the individual. We also report that the ^{14}C levels of two teeth from the same individual disclosed the order in which the teeth were formed, provided that their enamel formation times were sufficiently different. This can help to differentiate between teeth formed during the rising and falling part of the bomb-curve. However, using one single tooth, we also show that ^{14}C analysis of both enamel and root can accomplish the differentiation. This confirms the results that Cook *et al.* (2006) reported by analysis of ^{14}C in enamel and root collagen (23), but shows that extraction of the collagen constitutes an unnecessary step. Recently, Kondo-Nakamura *et al.* (2010) described a different approach, using separate analysis of the occlusal and cervical part of the enamel of the same tooth (24). However, they only describe such analysis of two teeth, and the difference in ^{14}C levels in the occlusal and cervical samples were very small. It is therefore more reliable to use either two teeth with sufficiently different enamel formation times or the enamel and root from the same tooth for ^{14}C analysis to determine if the tooth has been formed during the rising or falling part of the bomb-curve.

By analyzing teeth formed before the bomb spike we could show that there is some turnover of carbon in the root after completed formation. Since we only measured the mineral fraction in the roots, the incorporation of ^{14}C at adult ages in these roots shows that this component of the dentin and/or cementum continue to remodel, albeit at a slow pace given the relatively low ^{14}C levels found. From Table 2 it can further be appreciated that the teeth formed during the rising part of the curve (except Case 16) have higher levels of ^{14}C in the root than in the enamel, implying that if a continuous turnover were significant, the incorporation of the more recent atmospheric levels would have produced lower ^{14}C levels in the roots. In Case 16, the actual enamel formation time was calculated to be 1961.3, i.e. shortly before the atmospheric ^{14}C maximum. Given the long time period during which the roots are developed (completed about 6-7 years after the enamel (26)), most of the

incorporation of ^{14}C would have occurred during the falling part of the curve, explaining a lower ^{14}C level in the root than in the enamel.

The principle of estimating a person's date of birth from ^{14}C analysis of teeth has hitherto used reference data on tooth formation, in turn based on repeated radiographs of children (25). However, we have now compiled sufficient ^{14}C results for most types of teeth to bypass this step, so therefore we provide a reference tabulation of average ^{14}C incorporation times in Table S2 to be used for future calculations. These estimates generally show a shorter time interval than the radiographic results, which either may be due to our choice of Nolla's stage 4.5 as an average of the development, or explained by incorporation into enamel of some of the carbon earlier than the average radiographic picture of tooth formation. The radiocarbon that we measure with AMS is that incorporated in the carbonate component of the hydroxyapatite, and how this relates to the degree of radio-opacity is not known. Nonetheless, the use of crude ^{14}C levels in the enamel of teeth from subjects with a known DOB simplifies the calculation, and obviously provides a better precision of the DOB estimation of unknown individuals (Table S2).

Studies on hair samples from subjects in different US states (18) and in different countries (15) have shown that stable isotopes generate geographical signatures. Similarly, we have recently reported that such differences also can be seen in tooth enamel (16). In the present study we also show that ^{13}C levels in tooth enamel obviously vary within a limited geographical region such as the US and Canada. It seems likely that analysis of several stable isotopes in teeth and hair can help to determine both the earlier origin and more recent residence of an unknown dead body and thus facilitate identification work. Interestingly, the ^{13}C levels in teeth from subjects raised in Mexico are the highest we have recorded; even higher than in subjects from Chile and Uruguay. The most likely explanation is a high dietary intake of foods based on either corn or sugar cane, or both, since they represent so-called C4 plants that have double fixation steps in the photosynthesis system and fail to discriminate between ^{13}C and ^{12}C like C3 plants such as wheat, potato, rice and sugar beet (17, 27). The ^{13}C levels in tooth roots (Tables 4 and 5) were somewhat lower than in enamel from the same teeth and showed a higher variation within restricted regions, making it less reliable than enamel levels to determine geographic origin of the person. This might be explained by a more traditional diet with locally typical basic nutritional food sources during childhood than the diet during adolescence.

The ^{18}O levels in tooth roots were lowest in the northwest of the United States, which is in accordance with the map of levels in tap water previously reported (19), but showed somewhat variable levels in other parts of the United States (Fig. 1). The ^{18}O levels in body tissues may also be dependent on food intake with a variable level of this isotope, given that a substantial part of the diet is composed of water. Such contribution to the ^{18}O levels in tooth enamel may explain the smaller variation in teeth as compared to the variation in tap water, since the different components of diet may come from places at a distance from the residence - but perhaps not sufficiently remote to make a difference regarding ^{13}C .

The DNA analyses indicated the correct sex in all samples analyzed. The sex determination is important to separate possible matches from missing persons lists, but also for determining the year of birth of the person, since this calculation is based on tooth formation estimates and the tooth formation times are somewhat different between sexes for several types of teeth. DNA analysis are very swift and can be performed at short notice by many labs worldwide. The consistent results using two different methods for sex determination in select cases support the notion that these analyses are robust. In addition, since both the Identifiler™ kit and the Powerplex® Y System provide a specific profile of the individual, such an analysis will often be needed anyway if antemortem radiographs cannot be retrieved and it seems practical and more economical to perform this analysis sooner than later. The Doe network database includes a large number of unidentified persons and different identification methods may be used to compare the characteristics of the deceased with antemortem materials from the deceased or with samples from relatives. The review of this database (Table 1) revealed that the estimates of age of the subject show a great variation and that the sex was sometimes unknown. The analysis of bomb-pulse derived ^{14}C provides an accurate estimate of the year of birth of the person, but only the age at death when the year of death is known. We have previously shown that addition of established methods to determine the age at death, such as aspartic acid racemization, to the year of birth as estimated by ^{14}C measurement of teeth can be used to estimate the year of death of the person (5).

In conclusion, from one single tooth, the year of birth can be estimated, a clue to geographical origin can be obtained, sex can be determined and analysis of ^{14}C in both enamel and root can differentiate between tooth formation during rising and

falling part of the bomb-curve. We have also previously reported that a small portion of a tooth can be used for aspartic acid racemization to estimate the age at death of a person, which will allow for a calculation of the year of death, an important factor when investigating a skeletonized body.

In conclusion, a better characterization of unknown dead victims using these analyses can improve the identification work, whether it concerns a suspect homicide case or victims of a mass disaster.

Material and methods

Collection of teeth

In total, dentists in Mexico and the United States and Canada collected 66 teeth. The teeth were either extracted for orthodontic purposes or removed due to periodontal problems. Patients' permission to use the teeth for research instead of discarding them was obtained in all cases and the study was approved by the Regional Ethics Committee at the Karolinska Institute (No 2010/314-31/3). The tooth number, the year and month of birth, the sex, the area where the person was raised, and the date of extraction were noted and each person and tooth were given a code. All teeth were shipped to Karolinska Institute for preparation. A control determination of the tooth number was performed blindly with the assistance of three independent forensic odontologists.

Tooth preparation

Teeth were divided by cutting away the crown of the tooth from the root at the cervical line. To isolate the enamel, the excised crown was immersed in 10 N NaOH and placed in a water-bath sonicator overnight (Branson 150) at a maximum temperature of 70⁰C. Approximately every 24 hrs the NaOH was replaced and the softened non-enamel structures were removed by mechanical treatment using a dental drill and blunt dissection. Next, the enamel was washed in DDH₂O, re-submersed in 10N NaOH and returned to the sonicator. This procedure was repeated for 3-5 days until all dentin and soft structures were completely removed. The enamel was then rinsed several times in DDH₂O and dried at 65⁰C overnight. The enamel was weighed and kept sealed in a glass tube until pre-preparation for accelerator mass spectrometric (AMS) analysis. Roots were vertically divided to produce a somewhat

larger piece for isotope analysis and a smaller piece for DNA analysis. The roots were then rinsed in DDH₂O and no efforts were made to clean the root canal or to remove cementum.

Pre-treatment for AMS and stable isotope mass spectrometry

Aliquots of the enamel samples were placed in culture tubes for pre-treatment to remove the surface carbon that may have contaminated the enamel between formation and analysis. Aliquots of 80-150 mg were used to get full sized samples for ¹⁴C analysis. Enamel samples were immersed in 1.0 N HCl at room temperature for 1.5 hrs, rinsed 3 times with DDH₂O and placed on a heating block at 95⁰C to dry overnight. The acid pre-treatment was designed to etch the outer surface of the enamel that was exposed to the harsh base environment earlier without dissolving too much of the enamel. Each dried enamel sample was broken into 5-10 pieces and placed in an individual single-use reactor. Enamel splits in individual reaction chambers were evacuated, heated and acidified with concentrated orthophosphoric acid at 90⁰C to hydrolyze all mineralized carbon to CO₂. Roots were treated with 2% NaOCl for 24 h, then rinsed 10 times with DI water, followed by treatment with 0.1 N HCl for 30 min, again rinsed 10 times with DI water and finally dried on heating block overnight. 120 mg of the roots were then pulverized, and 1-2 mg of the powder was used for stable isotope analysis of ¹³C and ¹⁸O. Stable oxygen and carbon isotope ratios were determined by reaction of the powdered aliquot with supersaturated orthophosphoric acid at 90⁰C in an Isocarb common acid bath autocarbonate device attached to a Fisons Optima isotope ratio mass spectrometer. This method liberates carbonate so only the mineral fraction of the roots were used for stable isotope analysis.

The remainder of the powdered roots were used for AMS analysis of ¹⁴C and underwent similar processing as enamel after being placed in individual reactors. The evolved CO₂ from roots and enamel was purified, trapped, and reduced to graphite in the presence of iron catalyst in individual reactors (28, 29). The CO₂ from large enamel samples was split prior to graphitization and δ¹³C measured by stable isotope ratio mass spectrometry. Background values were controlled by consistently following procedures, frequently baking sample tubes, periodically cleaning rigs, and maintaining a clean lab (30).

Accelerator mass spectrometry analysis

Graphite targets were measured using the 10-MV HVEE FN-class tandem electrostatic AMS system at the Center for Accelerator Mass Spectrometry at Lawrence Livermore National Laboratory (LLNL). The operation is similar to that used when performing high-precision measurements of 18,000 year old turbidities used as secondary standards (31). The system employs a LLNL designed high-output negative ion solid graphite Cs-sputter source (32) which emits 250-350 μg of ^{12}C - from a full-sized sample, corresponding to approx. 900 ^{14}C counts per second from a contemporary sample. The FN AMS system routinely achieves 15 % total system efficiency for C analyzing $^{14}\text{C}^{4+}$ in the detector (33). Details on the design of the LLNL AMS system and its operation can be found in the literature (31-34). Enamel samples are usually full sized and contemporary, so analysis times are relatively rapid, generally less than 5 minutes. The enamel samples are measured for 30,000 ^{14}C counts per cycle for 5-7 cycle repetitions and achieve standard deviations of 0.3-0.8 %.

Corrections for background contamination introduced during AMS sample preparation are made by establishing the contributions from contemporary and fossil carbon, following the procedures of Brown and Southon (35). All data are normalized using six identically prepared NIST SRM 4990B (Oxalic Acid I) primary standards. NIST SRM 4990C (Oxalic Acid II), IAEA-C6 (36), and TIRI (37) wood are used as secondary standards and quality controls to monitor spectrometer performance. The ratio of NIST SRM 4990C to NIST SRM 4990B (Oxalic Acid II/Oxalic Acid I) measured between February 2005 and March 2012 on 31 different sample wheels containing enamel samples had an average value of 1.291 ± 0.003 (1 SD), in agreement with the certified value of 1.293 ± 0.001 . ^{14}C -free calcite serves as background material for processing the enamel samples. The enamel samples are organized in groups of 10-14 unknowns bracketed by primary standards with one primary standard in the middle of the group. The secondary standards, primary standards and group of unknowns are measured consecutively as a cycle. Upon completion of a cycle the set of primary standards, secondary standards and unknown samples are measured again until desired precision is achieved. A typical group of 14 enamel samples is measured completely in 2-3 h. The measurement error is

determined for each sample and generally ranges between $\pm 0.2 - 0.8 \%$ (1 SD). All ^{14}C data are reported using the $F^{14}\text{C}$ fraction modern nomenclature developed for post-bomb data (38). $F^{14}\text{C}$ is a concentration unit ($^{14}\text{C}/\text{C}$) denoting enrichment or depletion of ^{14}C relative to oxalic acid standard normalized for isotope fractionation. Data are also reported as decay corrected $\Delta^{14}\text{C}$ following the nomenclature of Stuiver and Pollach (39). $\Delta^{14}\text{C}$ was calculated using the equation:

$$\Delta^{14}\text{C} = 1000 * ((F^{14}\text{C} * \exp[\lambda*(1950 - y)] - 1))$$

where $\lambda = 1/8267 \text{ yr}^{-1}$ and $y = \text{year of measurement after 1950 A.D.}$

From ^{14}C to year of birth

The average age at which enamel formation is completed for each specific tooth has been determined previously and is dependent on the specific tooth and sex of the person (25, 26). In cases where the sex is unknown, the average time for enamel completion for males and females can be calculated. However in this study the sex was known for all individuals and confirmed by DNA analysis. To estimate an individual's date of birth the ^{14}C concentration measured in the tooth enamel was compared with the Calibomb reference data (<http://calib.qub.ac.uk/CALIBomb>) or Levin and Kromer (10) to find out the year of enamel formation of the particular tooth. By subtracting the average formation time for each type of tooth (25), the individual's date of birth can be calculated.

If it is not obvious whether an individual is born before or after the peak of atmospheric ^{14}C levels resulting from the bomb tests, then two teeth with different enamel formation times can be analyzed – this will distinguish whether the ^{14}C measurement relates to the rising or falling part of the curve. In 14 cases, two or more teeth from the same individual were subjected to AMS analysis. In addition, roots from the same teeth were also analyzed. The root of a tooth will have been formed later than the enamel and thus analysis of one single tooth can also provide such information (23, 26). This principle is illustrated in Fig. 3.

DNA analysis

DNA was extracted from small fragments of the 30 roots of teeth according to a previously described method (40) for bone DNA extraction. In short: the roots were placed in 96 % ethanol for a few minutes and rinsed with 0.5 % Na-hypochlorite and

dried overnight at 56°C in open tubes. The roots were then ground to a fine powder in liquid N₂ in a Freezer/Mill 6850-115 (SPEX Certiprep, New Jersey). The grinding cycles were the same as for bone treatment, but grinding was shortened to only half a minute. DNA was extracted from the total amount of tooth powder using one phenol-chlorophorm and one chlorophorm extraction (41), concentrated on Centricon 30 columns (Millipore, Billerica, MA), purified using the Qiaguick Purification Kit (Qiagen, Hilden, Germany) and eluted in 45 µL buffer.

The DNA concentration was measured with a NanoDrop® 2000 (Thermo Fischer Scientific, Wilmington, USA). Duplicate aliquots of approximately 1 ng were used for the PCR amplifications according to the Identifiler™ protocol (Applied Biosystems, Foster City, CA) in a final volume of 10 µL. To verify the results in weak samples these were also amplified with a two-phase PCR protocol of 10 and 20 cycles as previously described (40). The DNA profiles, including amelogenin, the marker for the determination of gender (42), were analyzed by capillary electrophoreses in an ABI3100 Genetic Analyzer (Applied Biosystems) and evaluated using GeneMapper ID 3.2 (Applied Biosystems). In select cases, the results of the amelogenin gene analysis was confirmed by analysis of Y chromosome specific STR markers using Powerplex® Y System (Promega Corp., Madison, WI), carried out according to the manufacturer's manual. The DNA extraction and amplification was performed using the same PCR procedure as outlined for the Identifiler™ protocol.

Statistical analysis

All results are given as means ± SD. Differences between two groups were examined for statistical significance with two-tailed Student's t test or with ANOVA, when appropriate. Significance was accepted at a *p* value of <.05.

Ethical considerations

All studies were approved by the Regional Ethical Committee in Stockholm, Sweden and patient permission obtained. All teeth were de-identified so that no new sensitive information of the individual subjects was produced.

Acknowledgments

All dentists procuring teeth are deeply acknowledged as well as the forensic

odontologists assisting in the confirmation of tooth type, in particular Irena Dawidson and Håkan Bengtsson. Support was provided by grants from the National Center for Research Resources (5P41RR013461), the National Institute of General Medical Sciences (8 P41 GM103483-14), and the National Institutes of Health and the Swedish Strategic Research Council (SSF). Work was performed in part under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

References

1. Ritz-Timme S, *et al.* (2000) Age estimation: The state of the art in relation to the specific demands of forensic practise. *Int J Legal Med* 113(3):129-136.
2. Helfman PM & Bada JL (1975) Aspartic acid racemization in tooth enamel from living humans. *Proc Natl Acad Sci U S A* 72(8):2891-2894.
3. Ohtani S & Yamamoto T (2005) Strategy for the estimation of chronological age using the aspartic acid racemization method with special reference to coefficient of correlation between D/L ratios and ages. *J Forensic Sci* 50(5):1020-1027.
4. Ohtani S & Yamamoto T (2010) Age estimation by amino acid racemization in human teeth. *J. Forensic. Sci.* 55(6):1630-3.
5. Alkass K, *et al.* (2010) Age estimation in forensic sciences: Application of combined aspartic acid racemization and radiocarbon analysis. *Mol Cell Proteomics* 9:1022-1030.
6. Spalding KL, Buchholz BA, Bergman LE, Druid H, & Frisen J (2005) Forensics: age written in teeth by nuclear tests. *Nature* 437(7057):333-334.
7. Libby WF, Berger R, Mead JF, Alexander GV, & Ross JF (1964) Replacement Rates for Human Tissue from Atmospheric Radiocarbon. *Science* 146:1170-1172.
8. De Vries H (1958) Atomic bomb effect: variation of radiocarbon in plants, shells, and snails in the past 4 years. *Science* 128(3318):250-251.
9. Hua Q & Barbetti M (2004) Review of tropospheric bomb ¹⁴C data for carbon cycle modeling and age calibration purposes. *Radiocarbon* 46(3):1273-1298.
10. Levin I & Kromer B (2004) The Tropospheric ¹⁴CO₂ Level in Mid-Latitudes of the Northern Hemisphere (1959–2003). *Radiocarbon* 46:1261-1272.
11. Nydal R & Lovseth K (1965) Distribution of radiocarbon from nuclear tests. *Nature* 206(988):1029-1031.
12. Spalding KL, Bhardwaj RD, Buchholz BA, Druid H, & Frisen J (2005) Retrospective birth dating of cells in humans. *Cell* 122(1):133-143.
13. Sweet DJ & Sweet CH (1995) DNA analysis of dental pulp to link incinerated remains of homicide victim to crime scene. *J Forensic Sci* 40(2):310-314.
14. Kelly JF, Atudorei V, Sharp ZD, & Finch DM (2002) Insights into Wilson's Warbler migration from analyses of hydrogen stable-isotope ratios. *Oecologia* 130(2):216-221.
15. Mutzel Rauch E, Lehn C, Peschel O, Holz S, & Rossmann A (2009) Assignment of unknown persons to their geographical origin by determination of stable isotopes in hair samples. *Int J Legal Med* 123(1):35-40.

16. Alkass K, Buchholz BA, Druid H, & Spalding KL (2011) Analysis of ¹⁴C and ¹³C in teeth provides precise birth dating and clues to geographical origin. *Forensic Sci Int.* 209(1-3):34-41.
17. Schoeller DA, Minigawa M, Slater R, & Kaplan IR (1986) Stable isotopes of carbon, nitrogen and hydrogen in the contemporary north american human food web. *Ecology of Food and Nutrition* 18:159-170.
18. Ehleringer JR, *et al.* (2008) Hydrogen and oxygen isotope ratios in human hair are related to geography. *Proc Natl Acad Sci U S A* 105(8):2788-2793.
19. Bowen GJ, Ehleringer JR, Chesson LA, Stange E, & Cerling TE (2007) Stable isotope ratios of tap water in the contiguous United States. *Water Resour Res* 43(3).
20. Gustafson G (1950) Age determination on teeth. *J Am Dent Assoc* 41(1):45-54.
21. Meinl A, Tangl S, Pernicka E, Fenes C, & Watzek G (2007) On the applicability of secondary dentin formation to radiological age estimation in young adults. *J Forensic Sci* 52(2):438-441.
22. Sullivan KM, Mannucci A, Kimpton CP, & Gill P (1993) A rapid and quantitative DNA sex test: fluorescence-based PCR analysis of X-Y homologous gene amelogenin. *Biotechniques* 15(4):636-638, 640-631.
23. Cook GT, Dunbar E, Black SM, & Xu S (2006) A preliminary assessment of age at death determination using the nuclear weapons testing C-14 activity of dentine and enamel. *Radiocarbon* 48(3):305-313.
24. Kondo-Nakamura M, Fukui K, Matsu'ura S, Kondo M, & Iwadate K (2010) Single tooth tells us the date of birth. *Int J Legal Med.*
25. Nolla CM (1960) The development of the permanent teeth. *J. Dental Child* 27.
26. Schour L & Massler M (1941) The development of the human dentition. *Am Dent Assoc* 28:1153.
27. O'Leary MH (1988) Carbon isotopes in photosynthesis. *BioScience* 38(5):328-336.
28. Santos GM, Southon JR, Druffel-Rodriguez K, Griffin S, & Mazon M (2004) Magnesium perchlorate as an alternative water trap in AMS graphite sample preparation: a report on sample preparation at KCCAMS at the University of California, Irvine. *Radiocarbon* 46:165-173.
29. Vogel JS, Southon JR, & Nelson DE (1987) Catalyst and Binder Effects in the Use of Filamentous Graphite for AMS. . *Nucl. Instrum. Methods Phys. Res. Sect. B* 29:50-56.
30. Zermeño P, *et al.* (2004) Prevention and removal of elevated radiocarbon contamination in the LLNL/CAMS natural radiocarbon sample preparation laboratory. *Nucl. Instrum. Methods Phys. Res. Sect. B* 223:293-297.
31. Guilderson TP, Southon JR, & Brown TA (2003) High-precision AMS C-14 results on TIRI/FIRI turbidite. *Radiocarbon* 45:75-81.
32. Davis JC, *et al.* (1990) LLNL/UC AMS Facility and Research Program. *Nucl. Instrum. Methods Phys. Res. Sect. B* 52:269-274.
33. Fallon SJ, Guilderson TP, & Brown TA (2007) CAMS/LLNL ion source efficiency revisited. *Nucl. Instrum. Methods Phys. Res. Sect. B* 259:106-110.
34. Southon JR & Roberts ML (2000) Ten years of sourcery at CAMS/LLNL – Evolution of a Cs ion source. *Nucl. Instrum. Methods Phys. Res. Sect. B* 172:257-261.


35. Brown TA & Southon JR (1997) Corrections for contamination background in AMS 14C measurements. *Nucl. Instrum. Methods Phys. Res. Sect. B* 123:208-213.
36. Rozanski K, *et al.* (1992) The IAEA C-14 intercomparison exercise 1990. *Radiocarbon* 34:506-519.
37. Scott EM (2003) The Third International Radiocarbon Intercomparison (TIRI) and the Fourth International Radiocarbon (FIR) - 1999-2002 - Results, analysis and conclusions *Radiocarbon* 45:293-408.
38. Reimer PJ, Brown TA, & Reimer RW (2004) Discussion: Reporting and calibration of post-bomb C-14 data. *Radiocarbon* 46:1299-1304.
39. Stuiver M & Pollach HA (1977) Discussion. Reporting of ¹⁴C data. *Radiocarbon* 19(3):355-363.
40. Winskog C, Nilsson H, Montelius K, & Lindblom B (2010) The use of commercial alcohol products to sterilize bones prior to DNA sampling. *Forensic Sci Med Pathol* 6(2):127-129.
41. Rainio J, *et al.* (2001) Forensic investigations in Kosovo: experiences of the European Union Forensic Expert Team. *J Clin Forensic Med* 8(4):218-221.
42. Nakahori Y, Takenaka O, & Nakagome Y (1991) A human X-Y homologous region encodes "amelogenin". *Genomics* 9(2):264-269.




Figure legends

Fig. 1. Map of ^{18}O in drinking water across the United States. The encircled values represent the average levels in tooth roots from subjects raised in the different states. Values from subjects from British Columbia and Washington are merged. The map is reproduced from (19) with permission by the publisher (American Geophysical Union, www.agu.org).

Fig. 2. Example of a part of a DNA profile obtained from analysis of a part of a tooth root. The IdentifilerTM kit is widely used and includes the profile of the amelogenin gene, here showing a result from a male subject (bottom panel).

Fig. 3. The principle for age estimation by analyzing ^{14}C in dental enamel. Horizontal green line represents a radiocarbon value crossing the ^{14}C bomb curve at two positions. By subtracting the average formation time for the particular tooth, the DOB of the person can be calculated (dashed green vertical line, next to actual DOB – solid green vertical line). However, if it is unclear whether the value matches the rising or falling part of the bomb curve, teeth with different laydown times can be analyzed (red marks), and thus indicate the correct part of the bomb curve.








Table 1. Panorama of the unknown bodies registered in the Doe Network database April 2011.

No. of Cases	Male (%)	Asian (%)	Black (%)	Hisp (%)	White (%)	N/A (%)	Avg. low¹	Avg. high¹	Error range²	>1 yr³ (%)
1229	66.7	2.6	51.7	6.2	20.6	18.9	29.1	43.0	15.0	49.0

¹The average minimum and maximum estimated age of the dead body.

²The average possible age range for all cases, where such an estimation is provided.

³This figure represents the number of cases that still are unidentified after more than a year.

Table 2. Overall results of ¹⁴C dating of teeth collected in North America.

Case No.	Sex	Tooth No.	Enamel formation time (yrs) ¹	Raised in / collected in	Radiocarbon analysis		Teeth	
					Fraction modern	±	Actual tooth DOB ¹	Estimated tooth DOB
1	M	41	2.5	BC	1.5410	0.0067	1965.3	1963.1
2	F	42	2.8	BC	0.9986	0.0038	1918.4	Pre-bomb
3	M	15	6.6	BC	1.5361	0.0045	1974.0	1970.1
4	M	44	5.1	BC	1.1130	0.0042	2001.7	1996,3
5	F	36	2.3	BC	1.3645	0.0051	1972.6	1967.4
6	M	22	4.0	BC	1.6698	0.0072	1966.2	1966.8
7	M	38	13.0	WA	1.1357	0.0040	1995.3	1992.5
	M	48	13.0	WA	1.1323	0.0042	1995.3	1993.1
8	F	27	5.8	WA	1.6459	0.0059	1968.6	1967.2
	F	45	5.7	WA	1.6356	0.0057	1968.5	1967.4
9	M	17	6.5	WA	1.1820	0.0042	1989.8	1987.5
10	M	47	6.5	WA	1.3421	0.0047	1979.9	1977.5
11	F	24	4.9	WA	1.3585	0.0050	1964.9	1962.4
12	M	16	3.3	WA	1.4898	0.0053	1973.9	1971.8
	M	37	6.5	WA	1.3518	0.0048	1977.1	1976.8
13	M	15	6.6	WA	1.1170	0.0040	1998.4	1995.6
14	M	17	6.5	MT/WA	1.1646	0.0041	1991.0	1989.4
	M	18	12.6	MT/WA	1.1242	0.0060	1997.1	1994.3
	M	28	12.6	MT/WA	1.1118	0.0039	1997.1	1996.5
15	F	25	5.6	MT/WA	1.3675	0.0048	1975.4	1975.8
	F	26	3.0	MT/WA	1.4750	0.0052	1972.8	1972.2
16	F	26	3.0	UK/WA ²	1.1765	0.0046	1990.6	1988.0
17	M	45	6.5	MA	1.2773	0.0036	1964.2	1962.0
18	M	44	5.1	CT	1.0940	0.0039	2002.3	2000.0
	M	14	5.6	CT	1.0922	0.0031	2002.8	2000.0
19	M	14	5.6	CT	1.0971	0.0038	2001.9	1999.2
	M	34	5.1	CT	1.0872	0.0038	2001.4	2001.0
20	F	31	2.5	CA	1.0134	0.0036	1947.4	Pre-bomb
	F	33	4.1	CA	0.9920	0.0029	1949.0	Pre-bomb
21	M	15	6.6	CA	1.0496	0.0044	1959.7	1957.3
22	M	28	12.6	CA/WA	1.1462	0.0040	1992.8	1991.6
23	M	42	3.0	GA	1.2281	0.0038	1984.1	1983.3
24	F	23	4.7	AL	1.0171	0.0036	1958.1	1956.1
25	F	14	4.9	TX	1.1396	0.0040	1996.3	1992.1
	F	34	4.4	TX	1.1224	0.0032	1995.8	1994.6
26	M	14	5.6	TX	0.9948	0.0028	1945.9	Pre-bomb
	M	31	2.5	TX	0.9900	0.0032	1942.8	Pre-bomb
27	M	23	4.7	TX	1.1726	0.0036	1960.8	1958,8
28	M	36	2.4	TX	1.1019	0.0035	1994.4	1998.2
29	M	18	12.6	TX	1.3799	0.0038	1976.5	1975.4
	M	46	2.4	TX	1.4075	0.0041	1967.6	1962.7
30	M	12	4.0	TX	1.3087	0.0038	1964.1	1962.1
	M	13	4.7	TX	1.3502	0.0041	1964.8	1962.4
	M	14	5.6	TX	1.4744	0.0062	1965.7	1962.9

	M	22	4.0	TX	1.6172	0.0066	1964.1	1963.3
	M	23	4.7	TX	1.7399	0.0267	1964.8	1965.8
	M	26	3.3	TX	1.2254	0.0046	1963.4	1961.1
	M	31	2.5	TX	1.2791	0.0073	1962.6	1962.0
	M	32	3.0	TX	1.6571	0.1693	1963.1	1963.3
	M	37	6.5	TX	1.6735	0.0056	1966.6	1966.8
	M	41	2.5	TX	1.4131	0.0059	1962.6	1962.6
	M	42	3.0	TX	1.5051	0.1504	1963.1	1962.8
	M	43	4.3	TX	1.4475	0.0062	1964.4	1962.8
	M	44	5.1	TX	1.6666	0.0067	1965.2	1966.4
	M	47	6.5	TX	1.6945	0.0060	1966.6	1966.5
31	M	11	3.2	TX	1.3500	0.0048	1979.6	1977.0
32	F	46	2.4	Mexico	0.9849	0.0035	1946.9	Pre-bomb
33	M	23	4.7	Mexico	1.0280	0.0040	1956.8	1956.0
34	M	26	3.3	Mexico	1.2163	0.0045	1962.8	1960.0
35	F	46	2.3	Mexico	0.9945	0.0035	1945.7	Pre-bomb
36	M	37	6.5	Mexico	1.6573	0.0059	1961.3	1962.7
	M	36	2.4	Mexico	1.0738	0.0038	1957.2	1957.4
37	F	35	5.7	Mexico	0.9920	0.0033	1926.7	Pre-bomb
38	F	36	2.3	Mexico	0.9949	0.0034	1933.5	Pre-bomb
39	F	17	5.8	Mexico	1.5306	0.0054	1969.6	1970.1
	F	45	5.7	Mexico	1.5331	0.0058	1969.5	1970.2

¹Enamel formation time according to Nolla (25).

²Unclear at what age this subject moved to WA.

Table 3. ^{14}C levels in enamel and roots from the same teeth.

Case No.	Tooth No.	Actual enamel DOB	Enamel		Root	
			Radiocarbon analysis			
			Fraction Modern	±	Fraction modern	±
2	42	1918.4	0.9986	0.0038	1.0579	0.0037
35	46	1945.7	0.9945	0.0035	1.0586	0.0037
26	14	1945.9	0.9948	0.0028	1.0194	0.0036
32	46	1946.8	0.9849	0.0035	1.0794	0.0038
20	33	1949.0	0.9920	0.0029	1.0662	0.0038
33	23	1956.8	1.0280	0.0040	1.4106	0.0050
24	23	1957.2	1.0171	0.0036	1.3740	0.0048
21	15	1959.7	1.0496	0.0044	1.2139	0.0048
27	23	1960.8	1.1726	0.0036	1.4541	0.0051
36	37	1961.3	1.6573	0.0059	1.4043	0.0056
30	32	1963.1	1.6571	0.1693	1.4007	0.0051
30	42	1963.1	1.5051	0.1504	1.4084	0.0051
1	41	1965.3	1.5410	0.0067	1.4372	0.0051
30	37	1966.6	1.6735	0.0056	1.3785	0.0044
39	45	1969.5	1.5331	0.0058	1.2894	0.0047
3	15	1974	1.5361	0.0045	1.2656	0.0044
15	25	1975.4	1.3675	0.0048	1.1948	0.0047
29	18	1977.9	1.3799	0.0038	1.1660	0.0042
31	11	1979.6	1.3500	0.0048	1.1708	0.0039
9	17	1989.8	1.1820	0.0042	1.1127	0.0037
16	26	1990.6	1.1765	0.0046	1.0910	0.0038
14	17	1991	1.1646	0.0041	1.0891	0.0043
22	28	1992.8	1.1462	0.0040	1.0865	0.0038
25	14	1996.3	1.1396	0.0040	1.0737	0.0038
4	44	2001.7	1.1130	0.0042	1.0575	0.0037
19	14	2001.9	1.0971	0.0038	1.0478	0.0039
18	14	2002.8	1.0922	0.0031	1.0442	0.0037
28	18	2004.6	1.1158	0.0032	1.0867	0.0049

Table 4. $\delta^{13}\text{C}$ levels in teeth from individuals raised in different parts of North America.

Case No.	Sex	Tooth No	Enamel formation time (yrs) ¹	Raised in	$\delta^{13}\text{C}$ enamel	$\delta^{13}\text{C}$ roots
3	M	15	6.6	BC	-11.78	-13.40
4	M	44	5.1	BC	-11.48	-12.22
5	F	36	2.3	BC	-10.99	
6	M	22	4	BC	-10.00	
12	M	37	6.5	WA	-11.74	
12	M	16	3.3	WA	-11.50	
11	F	24	4.9	WA	-10.35	
13	M	15	6.6	WA	-10.33	
9	M	17	6.5	WA	-10.03	-12.31
7	M	48	13	WA	-9.45	
7	M	38	13	WA	-9.38	
14	M	17	6.5	MT	-11.27	-6.27
15	F	26	3	MT	-11.07	
14	M	28	12.6	MT	-11.03	
17	M	45	6.5	MA	-7.89	
19	M	14	5.6	CT	-10.84	-12.20
18	M	14	5.6	CT	-10.82	-10.41
19	M	34	5.1	CT	-10.56	
22	M	28	12.6	CA	-12.56	-12.28
21	M	15	6.6	CA	-10.51	-11.28
20	F	33	4.1	CA	-10.39	-11.95
23	M	42	3	GA	-8.98	
24	F	23	3.8	AL	-9.02	-10.91
28	M	36	2.4	TX	-10.19	
25	F	14	4.9	TX	-9.54	-10.97
30	M	26	3.3	TX	-9.20	
25	F	34	4.4	TX	-9.18	
30	M	12	4.0	TX	-9.04	
29	M	46	2.4	TX	-8.96	-10.65
29	M	18	12.6	TX	-8.85	
26	M	14	5.6	TX	-8.83	-10.17
27	M	23	4.7	TX	-8.41	-9.92
36	M	37	6.5	Mexico	-7.78	-6.32
39	F	45	5.7	Mexico	-6.37	-9.55
36	M	36	2.4	Mexico	-6.29	
33	M	23	4.7	Mexico	-6.23	-8.77
34	M	26	3.3	Mexico	-5.76	
35	F	46	2.3	Mexico	-5.71	-6.80
32	F	46	2.3	Mexico	-4.32	-10.60
37	F	35	5.7	Mexico	-0.54	
38	F	36	2.3	Mexico	0.11	

¹Enamel formation time according to Nolla (25).

Table 5. $\delta^{18}\text{O}$ levels in tooth roots.

Case No.	Tooth No.	Raised in / collected in	$d^{18}\text{O}$ Root
3	15	BC	-8.03
1	41	BC	-6.23
2	42	BC	-9.88
4	44	BC	-8.12
9	17	WA	-7.14
16	26	UK/WA ¹	-6.35
15	25	MT/WA	-8.17
14	17	MT/WA	-5.67
19	14	CT	-6.26
18	14	CT	-5.00
22	28	CA/WA	-7.87
20	33	CA	-5.61
22	18	CA/WA	-8.12
21	15	CA	-4.81
24	23	AL	-8.15
30	42	TX	-9.01
31	11	TX	-5.48
28	18	TX	-5.15
25	14	TX	-5.14
30	32	TX	-5.11
29	18	TX	-5.64
30	37	TX	-5.16
26	14	TX	-5.02
27	23	TX	-5.95
30	33	TX	-5.44
32	46	Mexico	-7.16
39	45	Mexico	-6.60
33	23	Mexico	-6.43
37	37	Mexico	-6.38
35	46	Mexico	-6.24
36	37	Mexico	-6.43

¹Unclear at what age this subject moved to WA.

Table 6. Reference information for calculating the date of birth of the person using enamel laydown time and ^{14}C incorporation time.

Tooth No.	Enamel formation time ¹	^{14}C incorporation time ²	Absolute average error using enamel formation time ¹	Absolute average error using ^{14}C incorporation time ²
11/21	3.2	2.2 ± 1.4	1.5 ± 0.8	1.0 ± 0.9
12/22	3.9	3.8 ± 1.0	0.8 ± 0.6	0.9 ± 0.5
13/23	4.4	3.3 ± 1.3	1.5 ± 0.8	1.1 ± 0.7
14/24	5.2	3.6 ± 1.2	1.7 ± 1.1	1.0 ± 0.7
15/25	6.2	4.8 ± 1.3	1.5 ± 1.1	1.1 ± 0.7
16/26	3.2	1.5 ± 1.2	1.8 ± 1.0	1.0 ± 0.5
17/27	6.1	5.1 ± 1.0	1.2 ± 0.6	0.8 ± 0.6
18/28	12.1	11.5 ± 1.9	1.6 ± 1.1	1.4 ± 1.2
31/41	2.5	1.9 ± 0.8	0.6 ± 0.7	0.6 ± 0.5
32/42	3.0	3.2 ± 2.5	1.4 ± 1.9	1.5 ± 1.8
33/43	4.3	3.6 ± 0.9	0.8 ± 0.7	0.7 ± 0.2
34/44	4.9	4.1 ± 1.7	1.4 ± 1.2	1.1 ± 1.2
35/45	6.0	5.3 ± 0.8	0.8 ± 0.7	0.7 ± 0.3
36/46	2.4	3.0 ± 3.0	2.4 ± 1.7	2.2 ± 1.9
37/47	6.2	6.2 ± 1.6	1.2 ± 0.9	1.1 ± 1.1
38/48	12.6	11.0 ± 1.9	1.5 ± 1.1	1.1 ± 1.0
		Total.	1.4 ± 0.4	1.1 ± 0.5

¹Enamel formation time according to Nolla (25).

²The time interval used for error calculation does not include lag of ^{14}C in the food chain.

Supporting information.

Table S1 shows detailed data supporting Table 2. The $^{14}\text{C}/\text{C}$ ratios are corrected for isotope fractionation and expressed in both fraction modern, $F^{14}\text{C}$ (1) and decay corrected $\Delta^{14}\text{C}$ (2) nomenclatures. Whenever $\delta^{13}\text{C}$ could not be analyzed due to small sample size, the $\delta^{13}\text{C}$ was estimated from the average of teeth from that region with an uncertainty of $\pm 2\%$. Table S1 includes several teeth from some subjects. However, since the intra-individual results show similar variation as the inter-individual variation, their inclusion will not affect the overall error much.

Table S2 constitutes a compilation of the ^{14}C birth dating of tooth enamel included in this publication as well as such data from the three previous studies published by our laboratory (3-5). Only teeth prepared using the standard procedure – immersion in concentrated NaOH during sonication – are included. Cryo-cracking of tooth crowns and dissolution of whole crown are more rapid techniques, and might give similar precision. However, since the number of samples prepared by these methods are limited, we only present results from samples, in which the enamel has been isolated by the standard procedure. The table compares the precision of ^{14}C DOB estimation using reference information on enamel formation (6) with DOB estimation using the average ^{14}C incorporation time. It should be kept in mind that the ^{14}C levels in tooth enamel is not exactly the same as the contemporary atmospheric levels since there is a lag between atmosphere and human as carbon moves up the food chain (7, 8). Therefore the ^{14}C incorporation times are not an exact indicator of enamel formation time.

References

1. Reimer PJ, Brown TA, & Reimer RW (2004) Discussion: Reporting and calibration of post-bomb C-14 data. *Radiocarbon* 46(3):1299-1304.
2. Stuiver M, Polach, H. A. (1977) Discussion: reporting of 14C data. *Radiocarbon* 19:355-363.
3. Alkass K, Buchholz BA, Druid H, & Spalding KL (2011) Analysis of 14C and 13C in teeth provides precise birth dating and clues to geographical origin. *Forensic Sci Int* 209(1-3):34-41.
4. Alkass K, *et al.* (2010) Age estimation in forensic sciences: application of combined aspartic acid racemization and radiocarbon analysis. *Molecular & Cellular Proteomics* 9(5):1022-1030.
5. Spalding KL, Buchholz BA, Bergman LE, Druid H, & Frisen J (2005) Forensics: age written in teeth by nuclear tests. *Nature* 437(7057):333-334.
6. Nolla CM (1960) The development of the permanent teeth. *J. Dental Child* 27.
7. Libby WF, Berger R, Mead JF, Alexander GV, & Ross JF (1964) Replacement Rates for Human Tissue from Atmospheric Radiocarbon. *Science* 146:1170-1172.
8. Nydal R & Lovseth K (1965) Distribution of Radiocarbon from Nuclear Tests. *Nature* 206(4988):1029-1031.

Table S1. Background data and all $\delta^{14}\text{C}$ and $\delta^{13}\text{C}$ results for enamel.

Case No.	Sex	Tooth No.	Enamel formation time (yrs) ¹	Raised in / collected in	$\delta^{13}\text{C}^2$	$\delta^{14}\text{C}$	\pm	Fraction modern	\pm	Actual tooth DOB	Estimated tooth DOB	Error in yrs (2σ)	Actual DOB	Estimated DOB	Error	Absolute error
1	M	41	2.5	BC	-10.00	529.9	6.7	1.5410	0.0067	1965.3	1963.1	1.0	1962.8	1960.6	-2.2	2.2
2	F	42	2.8	BC	-11.00	-8.6	3.8	0.9986	0.0038	1918.4	Pre-bomb	Pre-bomb	1915.6	Pre-bomb	Pre-bomb	Pre-bomb
3	M	15	6.6	BC	-11.78	525.0	4.5	1.5361	0.0045	1974.0	1970.1	0.1	1967.4	1963.5	-3.9	3.9
4	M	44	5.1	BC	-11.48	104.9	4.2	1.1130	0.0042	2001.7	1996.3	2.1	1996.6	1991.2	-5.4	5.4
5	F	36	2.3	BC	-10.99	354.3	5.1	1.3645	0.0051	1972.6	1967.4	1.0	1970.4	1974.2	3.8	3.8
6	M	22	4.0	BC	-10.00	657.4	7.2	1.6698	0.0072	1966.2	1966.8	0.4	1962.2	1962.9	0.7	0.7
7	M	38	13.0	WA	-10.00	127.3	4.0	1.1357	0.0040	1995.3	1992.5	1.7	1982.3	1979.5	-2.8	2.8
7	M	48	13.0	WA	-10.00	124.0	4.2	1.1323	0.0042	1995.3	1993.1	1.9	1982.3	1980.1	-2.2	2.2
8	F	27	5.8	WA	-10.00	633.8	5.9	1.6459	0.0059	1968.6	1967.2	0.3	1962.8	1961.4	-1.4	1.4
8	F	45	5.7	WA	-10.00	623.6	5.7	1.6356	0.0057	1968.5	1967.4	0.3	1962.8	1961.7	-1.1	1.1
9	M	17	6.5	WA	-10.00	173.3	4.2	1.1820	0.0042	1989.8	1987.5	1.4	1983.3	1981.0	-2.3	2.3
10	M	47	6.5	WA	-10.00	332.3	4.7	1.3421	0.0047	1979.9	1977.5	1.0	1973.4	1971.0	-2.4	2.4
11	F	24	4.9	WA	-10.00	348.5	5.0	1.3585	0.0050	1964.9	1962.4	0.2	1960.0	1957.5	-2.5	2.5
12	M	16	3.3	WA	-10.00	478.8	5.3	1.4898	0.0053	1973.9	1971.8	0.5	1970.6	1968.5	-2.1	2.1
12	M	37	6.5	WA	-10.00	341.9	4.8	1.3518	0.0048	1977.1	1976.8	1.0	1970.6	1970.3	-0.3	0.3
13	M	15	6.6	WA	-10.00	108.8	4.0	1.1170	0.0040	1998.4	1995.6	1.9	1991.8	1989.0	-2.8	2.8
14	M	17	6.5	MT	-10.00	156.1	4.1	1.1646	0.0041	1991.0	1989.4	0.5	1984.5	1982.9	-1.6	1.6
14	M	18	12.6	MT	-10.00	115.9	6.0	1.1242	0.0060	1997.1	1994.3	2.4	1984.5	1981.7	-2.8	2.8
14	M	28	12.6	MT	-10.00	103.6	3.9	1.1118	0.0039	1997.1	1996.5	2.1	1984.5	1983.9	-0.6	0.6
15	F	25	5.6	MT	-10.00	357.5	4.8	1.3675	0.0048	1975.4	1975.8	0.8	1969.8	1970.2	0.4	0.4
15	F	26	3.0	MT	-10.00	464.1	5.2	1.4750	0.0052	1972.8	1972.2	0.4	1969.8	1969.2	-0.6	0.6
17	M	45	6.5	MA	-7.89	267.7	3.6	1.2773	0.0036	1964.2	1962.0	0.2	1957.7	1956.0	-1.7	1.7
18	M	14	5.6	CT	-10.82	84.3	3.1	1.0922	0.0031	2002.8	2000.0	1.8	1997.2	1994.4	-2.8	2.8
18	M	44	5.1	CT	-10.00	86.1	3.9	1.0940	0.0039	2002.3	2000.0	1.9	1997.2	1994.9	-2.3	2.3
19	M	14	5.6	CT	-10.84	89.2	3.8	1.0971	0.0038	2001.9	1999.2	1.9	1996.3	1993.6	-2.7	2.7
19	M	34	5.1	CT	-10.56	79.4	3.8	1.0872	0.0038	2001.4	2001.0	1.1	1996.3	1995.9	-0.4	0.4

16	F	26	3.0	WA ³	-10.00	167.9	4.6	1.1765	0.0046	1990.6	1988.0	1.5	1987.6	1985.0	-2.6	2.6
20	F	31	2.5	CA	-10.00	6.1	3.6	1.0134	0.0036	1947.4	Pre-bomb	Pre-bomb	1944.9	Pre-bomb	Pre-bomb	Pre-bomb
20	F	33	4.1	CA	-10.39	-15.2	2.9	0.9920	0.0029	1949.0	Pre-bomb	Pre-bomb	1944.9	Pre-bomb	Pre-bomb	Pre-bomb
21	M	15	6.6	CA	-10.51	42.0	4.4	1.0496	0.0044	1959.7	1957.3	0.5	1953.1	1950.7	-2.4	2.4
22	M	28	12.6	CA	-10.00	137.8	4.0	1.1462	0.0040	1992.8	1991.6	1.4	1980.2	1979.0	-1.2	1.2
23	M	42	3.0	GA	-8.98	218.9	3.8	1.2281	0.0038	1984.1	1983.3	0.8	1981.1	1980.3	-0.7	0.7
24	F	23	4.7	AL	-10.00	9.6	3.6	1.0171	0.0036	1958.1	1956.1	0.5	1953.4	1951.4	-2.0	2.0
25	F	14	4.9	TX	-9.54	131.3	4.0	1.1396	0.0040	1996.3	1992.1	1.5	1991.4	1987.2	-4.2	4.2
25	F	34	4.4	TX	-9.18	114.2	3.2	1.1224	0.0032	1995.8	1994.6	1.5	1991.4	1990.2	-1.2	1.2
26	M	14	5.6	TX	-8.83	-12.4	2.8	0.9948	0.0028	1945.9	Pre-bomb	Pre-bomb	1940.3	Pre-bomb	Pre-bomb	Pre-bomb
26	M	31	2.5	TX	-9.00	-17.2	3.2	0.9900	0.0032	1942.8	Pre-bomb	Pre-bomb	1940.3	Pre-bomb	Pre-bomb	Pre-bomb
27	M	23	4.7	TX	-8.41	164.1	3.6	1.1726	0.0036	1960.8	1958.8	0.5	1956.1	1954.1	-2.0	2.0
28	M	36	2.4	TX	-10.19	93.9	3.5	1.1019	0.0035	1994.4	1998.2	2.2	1992.0	1995.8	3.8	3.8
29	M	18	12.6	TX	-8.85	370.0	3.8	1.3799	0.0038	1976.5	1975.4	0.7	1965.3	1962.8	-2.5	2.5
29	M	46	2.4	TX	-8.96	397.3	4.1	1.4075	0.0041	1967.6	1962.7	0.2	1965.3	1960.3	-5.0	5.0
30	M	12	4.0	TX	-9.04	299.2	3.8	1.3087	0.0038	1964.1	1962.1	0.1	1960.1	1958.1	-2.0	2.0
30	M	13	4.7	TX	-9.00	340.4	4.1	1.3502	0.0041	1964.8	1962.4	0.2	1960.1	1957.7	-2.4	2.4
30	M	14	5.6	TX	-9.00	463.8	6.2	1.4744	0.0062	1965.7	1962.9	0.4	1960.1	1957.3	-2.8	2.8
30	M	22	4.0	TX	-9.00	605.5	6.6	1.6172	0.0066	1964.1	1963.3	0.3	1960.1	1959.3	-0.8	0.8
30	M	23	4.7	TX	-9.00	727.3	26.7	1.7399	0.0267	1964.8	1965.8	1.0	1960.1	1961.1	1.0	1.0
30	M	26	3.3	TX	-9.20	216.5	4.6	1.2254	0.0046	1963.4	1961.1	0.5	1960.1	1957.8	-2.3	2.3
30	M	31	2.5	TX	-9.00	269.9	7.3	1.2791	0.0073	1962.6	1962.0	0.4	1960.1	1959.5	-0.6	0.6
30	M	32	3.0	TX	-9.00	645.2	169.3	1.6571	0.1693	1963.1	1963.3	0.5	1960.1	1960.3	0.2	0.2
30	M	37	6.5	TX	-9.00	661.4	5.6	1.6735	0.0056	1966.6	1966.8	0.4	1960.1	1960.3	0.2	0.2
30	M	41	2.5	TX	-9.00	402.8	5.9	1.4131	0.0059	1962.6	1962.6	0.3	1960.1	1960.1	0.0	0.0
30	M	42	3.0	TX	-9.00	494.3	150.4	1.5051	0.1504	1963.1	1962.8	0.6	1960.1	1959.8	-0.3	0.3
30	M	43	4.3	TX	-9.00	437.1	6.2	1.4475	0.0062	1964.4	1962.8	0.2	1960.1	1958.5	-1.6	1.6
30	M	44	5.1	TX	-9.00	654.6	6.7	1.6666	0.0067	1965.2	1966.4	0.1	1960.1	1961.3	1.2	1.2
30	M	47	6.5	TX	-9.00	682.3	6.0	1.6945	0.0060	1966.6	1966.5	0.4	1960.1	1960.0	-0.1	0.1
31	M	11	3.2	TX	-10.00	340.3	4.8	1.3500	0.0048	1979.6	1977.0	1.0	1976.4	1973.8	-2.6	2.6
32	F	46	2.4	Mexico	-10.00	-22.3	3.5	0.9849	0.0035	1946.9	Pre-bomb	Pre-bomb	1944.5	Pre-bomb	Pre-bomb	Pre-bomb

33	M	23	4.7	Mexico	<i>-10.00</i>	20.4	4.0	1.0280	0.0040	1956.8	1956.0	0.5	1952.1	1951.3	-0.8	0.8
34	M	26	3.3	Mexico	<i>-10.00</i>	207.4	4.5	1.2163	0.0045	1962.8	1960.0	1.4	1959.5	1956.7	-2.8	2.8
35	F	46	2.3	Mexico	<i>-10.00</i>	-12.8	3.5	0.9945	0.0035	1945.7	Pre-bomb	Pre-bomb	1943.4	Pre-bomb	Pre-bomb	Pre-bomb
36	M	37	6.5	Mexico	<i>-10.00</i>	645.1	5.9	1.6573	0.0059	1961.3	1962.7	0.5	1954.8	1956.2	1.4	1.4
36	M	36	2.4	Mexico	<i>-10.00</i>	65.9	3.8	1.0738	0.0038	1957.2	1957.4	0.5	1954.8	1955.0	0.2	0.2
37	F	35	5.7	Mexico	<i>-10.00</i>	-15.3	3.3	0.9920	0.0033	1926.7	Pre-bomb	Pre-bomb	1921.0	Pre-bomb	Pre-bomb	Pre-bomb
38	F	36	2.3	Mexico	<i>-10.00</i>	-12.4	3.4	0.9949	0.0034	1933.5	Pre-bomb	Pre-bomb	1931.2	Pre-bomb	Pre-bomb	Pre-bomb
39	F	17	5.8	Mexico	<i>-10.00</i>	519.3	5.4	1.5306	0.0054	1969.6	1970.1	0.5	1963.8	1964.3	0.5	0.5
39	F	45	5.7	Mexico	<i>-10.00</i>	521.9	5.8	1.5331	0.0058	1969.5	1970.2	0.8	1963.8	1964.5	0.7	0.7

Average **1.9**
SD 1.3

¹Enamel formation time according to Nolla (25).

²Values in italics are estimates for each region.

³This person was born in England, UK, but moved to WA, unclear at what age.

Table S2. Compilation of DOB estimation using average ^{14}C incorporation time interval only and using both ^{14}C and enamel laydown data.

Sex	Country ¹	Tooth No.	Actual DOB of person	Enamel formation time (yrs) ²	F ¹⁴ C	+/-	Estimated tooth DOB using only ¹⁴ C data ³	Interval DOB - ¹⁴ C incorporation	Estimated tooth DOB using enamel formation time	Estimated DOB using only ¹⁴ C data	Error using enamel formation time	Absolute error using enamel formation time	Error using only ¹⁴ C data	Absolute error using only ¹⁴ C data
M	TX	11	1976.4	3.2	1.3500	0.0048	1977.0	0.6	1973.8	1974.8	-2.6	2.6	-1.6	1.6
F	n/a	11	1957.7	3.2	1.1310	0.0070	1959.8	2.1	1956.6	1957.6	-1.1	1.1	-0.1	0.1
M	Sweden	21	1982.0	3.2	1.2138	0.0031	1984.4	2.4	1981.2	1982.2	-0.8	0.8	0.2	0.2
M	Lebanon	21	1962.6	3.2	1.5237	0.0164	1963.0	0.4	1959.8	1960.8	-2.8	2.8	-1.8	1.8
F	n/a	21	1957.7	3.2	1.1420	0.0070	1959.8	2.1	1956.6	1957.6	-1.1	1.1	-0.1	0.1
F	Scand?	21	1970.4	3.2	1.3920	0.0060	1974.8	4.4	1971.6	1972.6	1.2	1.2	2.2	2.2
F	n/a	21	1975.0	3.2	1.3150	0.0060	1978.8	3.8	1975.6	1976.6	0.6	0.6	1.6	1.6
F	Poland	11	1957.0	3.2	1.1471	0.0045	1958.5	1.5	1955.3	1956.3	-1.7	1.7	-0.7	0.7
			Mean	3.2			Mean	2.2		Mean	-1.0	1.5	0.0	1.0
							SD	1.4		SD	1.4	0.8	1.4	0.9
M	TX	12	1960.1	4.0	1.3087	0.0038	1962.1	2.0	1958.2	1958.4	-1.9	1.9	-1.8	1.8
F	n/a	12	1990.5	3.7	1.1200	0.0040	1995.2	4.7	1991.3	1991.5	0.8	0.8	1.0	1.0
F	n/a	12	1975.0	3.7	1.2930	0.0060	1979.7	4.7	1975.8	1976.0	0.8	0.8	1.0	1.0
F	n/a	12	1971.8	3.7	1.4010	0.0080	1974.4	2.6	1970.5	1970.7	-1.3	1.3	-1.1	1.1
M	TX	22	1960.1	4.0	1.6172	0.0066	1963.3	3.2	1959.4	1959.6	-0.7	0.7	-0.5	0.5
F	n/a	22	1975.0	3.7	1.3100	0.0060	1979.0	4.0	1975.1	1975.3	0.1	0.1	0.3	0.3
M	Bosnia	22	1953.0	4.0	1.0471	0.0038	1957.2	4.2	1953.3	1953.5	0.3	0.3	0.5	0.5
M	BC	22	1962.2	4.0	1.6698	0.0072	1966.8	4.6	1962.9	1963.1	0.7	0.7	0.8	0.8
			Mean	3.9			Mean	3.8		Mean	-0.2	0.8	0.0	0.9

							SD	1.0		SD	1.0	0.6	1.0	0.5
F	Japan	13	1967.3	3.8	1.5434	0.0060	1970.1	2.8	1965.7	1966.8	-1.6	1.6	-0.5	0.5
F	n/a	13	1990.5	3.8	1.1200	0.0050	1995.3	4.8	1990.9	1992.0	0.4	0.4	1.5	1.5
M	TX	13	1960.1	4.7	1.3502	0.0041	1962.4	2.3	1958.0	1959.1	-2.1	2.1	-1.0	1.0
F	AL	23	1953.4	4.7	1.0171	0.0036	1956.1	2.7	1951.7	1952.8	-1.7	1.7	-0.6	0.6
M	Mexico	23	1952.1	4.7	1.0280	0.0040	1956.0	3.9	1951.6	1952.7	-0.5	0.5	0.6	0.6
M	TX	23	1956.1	4.7	1.1726	0.0036	1958.8	2.7	1954.4	1955.5	-1.7	1.7	-0.6	0.6
M	TX	23	1960.1	4.7	1.7399	0.0267	1965.8	5.7	1961.4	1962.5	1.3	1.3	2.4	2.4
F	Poland	13	1957.0	3.8	1.1403	0.0041	1958.4	1.4	1954.0	1955.1	-3.0	3.0	-1.9	1.9
			Mean	4.4			Mean	3.3		Mean	-1.1	1.5	0.0	1.1
							SD	1.3		SD	1.4	0.8	1.4	0.7
F	n/a	14	1991.5	4.9	1.1251	0.0072	1994.2	2.7	1989.0	1990.6	-2.5	2.5	-0.9	0.9
F	Sweden	14	1988.8	4.9	1.1533	0.0032	1990.6	1.8	1985.4	1987.0	-3.4	3.4	-1.8	1.8
F	Sweden	14	1991.3	4.9	1.1088	0.0040	1997.2	5.9	1992.0	1993.6	0.7	0.7	2.3	2.3
F	Sweden	14	1989.1	4.9	1.1390	0.0045	1992.2	3.1	1987.0	1988.6	-2.1	2.1	-0.5	0.5
F	Sweden	14	1991.1	4.9	1.1195	0.0041	1995.2	4.1	1990.0	1991.6	-1.1	1.1	0.5	0.5
F	Sweden	14	1991.5	4.9	1.1177	0.0041	1995.5	4.0	1990.3	1991.9	-1.2	1.2	0.4	0.4
F	Sweden	14	1988.3	4.9	1.1413	0.0037	1992.0	3.7	1986.8	1988.4	-1.5	1.5	0.1	0.1
F	TX	14	1991.4	4.9	1.1396	0.0040	1992.1	0.7	1986.9	1988.5	-4.5	4.5	-2.9	2.9
M	Chile	14	1967.8	5.6	1.4825	0.0057	1972.0	4.2	1966.8	1968.4	-1.0	1.0	0.6	0.6
M	Sweden	14	1993.2	5.6	1.1024	0.0042	1998.1	4.9	1992.9	1994.5	-0.3	0.3	1.3	1.3
M	Sweden	14	1990.9	5.6	1.1217	0.0043	1994.9	4.0	1989.7	1991.3	-1.2	1.2	0.4	0.4
M	Japan	14	1964.2	5.6	1.5797	0.0061	1968.4	4.2	1963.2	1964.8	-1.0	1.0	0.6	0.6
M	CT	14	1997.2	5.6	1.0922	0.0031	2000.0	2.8	1994.8	1996.4	-2.4	2.4	-0.8	0.8
M	CT	14	1996.3	5.6	1.0971	0.0038	1999.2	2.9	1994.0	1995.6	-2.3	2.3	-0.7	0.7

M	TX	14	1960.1	5.6	1.4744	0.0062	1962.9	2.8	1957.7	1959.3	-2.4	2.4	-0.8	0.8
M	Sweden	14	1990.8	5.6	1.1196	0.0043	1995.3	4.5	1990.1	1991.7	-0.7	0.7	0.9	0.9
F	WA	24	1960.0	4.9	1.3585	0.0050	1962.4	2.4	1957.2	1958.8	-2.8	2.8	-1.2	1.2
F	Sweden	24	1990.5	4.9	1.1215	0.0054	1994.9	4.4	1989.7	1991.3	-0.8	0.8	0.8	0.8
F	Sweden	24	1991.8	4.9	1.1133	0.0042	1996.2	4.4	1991.0	1992.6	-0.8	0.8	0.8	0.8
M	n/a	24	1992.1	5.6	1.1198	0.0050	1995.3	3.2	1990.1	1991.7	-2.0	2.0	-0.4	0.4
M	Scand?	24	1993.1	5.6	1.1030	0.0040	1998.0	4.9	1992.8	1994.4	-0.3	0.3	1.3	1.3
			Mean	5.2			Mean	3.6		Mean	-1.6	1.7	0.0	1.0
							SD	1.2		SD	1.2	1.1	1.2	0.7
F	N	15	1989.1	5.6	1.1301	0.0039	1993.4	4.3	1987.2	1988.6	-1.9	1.9	-0.5	0.5
F	Scand?	15	1988.6	5.6	1.1264	0.0043	1993.8	5.2	1987.6	1989.0	-1.0	1.0	0.4	0.4
F	Na	15	1989.5	5.6	1.1120	0.0040	1996.5	7.0	1990.3	1991.7	0.8	0.8	2.2	2.2
M	BC	15	1967.4	6.6	1.5361	0.0045	1970.1	2.7	1963.9	1965.3	-3.5	3.5	-2.1	2.1
M	CA	15	1953.1	6.6	1.0496	0.0044	1957.3	4.2	1951.1	1952.5	-2.0	2.0	-0.6	0.6
M	WA	15	1991.8	6.6	1.1170	0.0040	1995.6	3.8	1989.4	1990.8	-2.4	2.4	-1.0	1.0
M	Chile	15	1967.8	6.6	1.4785	0.0236	1972.1	4.3	1965.9	1967.3	-1.9	1.9	-0.5	0.5
M	Uruguay	25	1977.4	6.6	1.2275	0.0044	1983.4	6.0	1977.2	1978.6	-0.2	0.2	1.2	1.2
F	MT/WA	25	1969.8	5.6	1.3675	0.0048	1975.8	6.0	1969.6	1971.0	-0.2	0.2	1.2	1.2
			Mean	6.2			Mean	4.8		Mean	-1.4	1.5	0.0	1.1
							SD	1.3		SD	1.3	1.1	1.3	0.7
M	WA	16	1970.6	3.3	1.4898	0.0053	1971.8	1.2	1968.6	1970.3	-2.0	2.0	-0.3	0.3
M	TX	26	1960.1	3.3	1.2254	0.0046	1961.1	1.0	1957.9	1959.6	-2.2	2.2	-0.5	0.5
M	Chile	26	1967.8	3.3	1.5173	0.0108	1970.8	3.0	1967.6	1969.3	-0.2	0.2	1.5	1.5
M	Scand?	26	1956.5	3.3	1.0543	0.0036	1957.1	0.6	1953.9	1955.6	-2.6	2.6	-0.9	0.9
M	Mexico	26	1959.5	3.3	1.2163	0.0045	1960.0	0.5	1956.8	1958.5	-2.7	2.7	-1.0	1.0

F	Uruguay	26	1995.2	3.0	1.1137	0.0036	1996.1	0.9	1992.9	1994.6	-2.3	2.3	-0.6	0.6
F	England/WA ⁴	26	1987.6	3.0	1.1765	0.0046	1988.0	0.4	1984.8	1986.5	-2.8	2.8	-1.1	1.1
F	MT/WA	26	1969.8	3.0	1.4750	0.0052	1972.2	2.4	1969.0	1970.7	-0.8	0.8	0.9	0.9
F		16	1955.1	3.0	1.0089	0.0037	1958.7	3.6	1955.5	1957.2	0.4	0.4	2.1	2.1
			Mean	3.2			Mean	1.5		Mean	-1.7	1.8	0.0	1.0
							SD	1.2		SD	1.2	1.0	1.2	0.5
F	N	17	1972.2	5.8	1.3359	0.0042	1977.8	5.6	1971.7	1972.7	-0.5	0.5	0.5	0.5
F	Sweden	17	1993.4	5.8	1.1035	0.0036	1998.1	4.7	1992.0	1993.0	-1.4	1.4	-0.4	0.4
F	Sweden	17	1963.7	5.8	1.5653	0.0051	1969.4	5.7	1963.3	1964.3	-0.4	0.4	0.6	0.6
F	Scand?	17	1955.8	5.8	1.2160	0.0038	1960.0	4.2	1953.9	1954.9	-1.9	1.9	-0.9	0.9
F	Mexico	17	1963.8	5.8	1.5306	0.0054	1970.1	6.3	1964.0	1965.0	0.2	0.2	1.2	1.2
M	MT/WA	17	1984.5	6.5	1.1646	0.0041	1989.4	4.9	1983.3	1984.3	-1.2	1.2	-0.2	0.2
M	WA	17	1983.3	6.5	1.1820	0.0042	1987.5	4.2	1981.4	1982.4	-1.9	1.9	-0.9	0.9
M	Scand?	17	1952.3	6.5	1.0277	0.0026	1956.4	4.1	1950.3	1951.3	-2.0	2.0	-1.0	1.0
M	Iraq	27	1975.8	6.5	1.2706	0.0042	1980.8	5.0	1974.7	1975.7	-1.1	1.1	-0.1	0.1
M	Scand?	27	1989.6	6.5	1.1167	0.0038	1997.0	7.4	1990.9	1991.9	1.3	1.3	2.3	2.3
F	WA	27	1962.8	5.8	1.6459	0.0059	1967.2	4.4	1961.1	1962.1	-1.7	1.7	-0.7	0.7
			Mean	6.1			Mean	5.1		Mean	-1.0	1.2	0.0	0.8
							SD	1.0		SD	1.0	0.6	1.0	0.6
F	Scand?	18	1965.3	11.2	1.3065	0.0038	1979.2	13.9	1967.1	1967.7	1.8	1.8	2.4	2.4
M	Scand?	18	1962.6	12.6	1.4280	0.0045	1973.7	11.1	1961.6	1962.2	-1.0	1.0	-0.4	0.4
M	Scand?	18	1977.1	12.6	1.1284	0.0033	1993.6	16.5	1981.5	1982.1	4.4	4.4	5.0	5.0
M	MT/WA	18	1984.5	12.6	1.1242	0.0060	1994.3	9.8	1982.2	1982.8	-2.3	2.3	-1.7	1.7
M	TX	18	1965.3	12.6	1.3799	0.0038	1975.4	10.1	1963.3	1963.9	-2.0	2.0	-1.4	1.4

M	Chile	28	1967.8	12.6	1.3236	0.0317	1978.2	10.4	1966.1	1966.7	-1.7	1.7	-1.1	1.1
M	Morocco	28	1958.3	12.6	1.4386	0.0049	1973.4	15.1	1961.3	1961.9	3.0	3.0	3.6	3.6
M		28	1967.3	12.6	1.3210	0.0050	1978.4	11.1	1966.3	1966.9	-1.0	1.0	-0.4	0.4
M	Scand?	28	1985.7	12.6	1.1137	0.0031	1996.1	10.4	1984.0	1984.6	-1.7	1.7	-1.1	1.1
M	Scand?	28	1956.7	12.6	1.5728	0.0050	1968.6	11.9	1956.5	1957.1	-0.2	0.2	0.4	0.4
M	Scand?	28	1949.4	12.6	1.3612	0.0066	1962.4	13.0	1950.3	1950.9	0.9	0.9	1.5	1.5
M	CA/WA	28	1980.2	12.6	1.1462	0.0040	1991.6	11.4	1979.5	1980.1	-0.7	0.7	-0.1	0.1
M	MT/WA	28	1984.5	12.6	1.1118	0.0039	1996.5	12.0	1984.4	1985.0	-0.1	0.1	0.5	0.5
F	Scand?	28	1956.8	11.2	1.7039	0.0064	1966.4	9.6	1954.3	1954.9	-2.5	2.5	-1.9	1.9
F	Scand?	28	1969.3	11.2	1.3044	0.0044	1979.2	9.9	1967.1	1967.7	-2.2	2.2	-1.6	1.6
F	Scand?	28	1968.8	11.2	1.2973	0.0047	1979.6	10.8	1967.5	1968.1	-1.3	1.3	-0.7	0.7
F	Scand?	28	1977.2	11.2	1.1757	0.0039	1988.0	10.8	1975.9	1976.5	-1.3	1.3	-0.7	0.7
F	Uruguay	28	1985.6	11.2	1.1198	0.0032	1995.1	9.5	1983.0	1983.6	-2.6	2.6	-2.0	2.0
F	Lebanon	28	1985.1	11.2	1.1084	0.0051	1996.9	11.8	1984.8	1985.4	-0.3	0.3	0.3	0.3
			Mean	12.1			Mean	11.5		Mean	-0.6	1.6	0.0	1.4
							SD	1.9		SD	1.9	1.1	1.9	1.2
M	Scand?	31	1958.2	2.5	1.2077	0.0047	1960.4	2.2	1957.9	1958.5	-0.3	0.3	0.3	0.3
M	TX	31	1960.1	2.5	1.2791	0.0073	1962.0	1.9	1959.5	1960.1	-0.6	0.6	0.0	0.0
M	BC	41	1962.8	2.5	1.5410	0.0067	1963.1	0.3	1960.6	1961.2	-2.2	2.2	-1.6	1.6
M	TX	41	1960.1	2.5	1.4131	0.0059	1962.6	2.5	1960.1	1960.7	0.0	0.0	0.6	0.6
F	Scand?	41	1970.2	2.5	1.4480	0.0060	1973.0	2.8	1970.5	1971.1	0.3	0.3	0.9	0.9
F	Scand?	41	1990.5	2.5	1.1340	0.0050	1992.8	2.3	1990.3	1990.9	-0.2	0.2	0.4	0.4
M		41	1973.0	2.5	1.3982	0.0052	1974.6	1.6	1972.1	1972.7	-0.9	0.9	-0.3	0.3
			Mean	2.5			Mean	1.9		Mean	-0.6	0.6	0.0	0.6
							SD	0.8		SD	0.8	0.7	0.8	0.5

M	TX	32	1960.1	3.0	1.6571	0.1693	1963.3	3.2	1960.3	1960.1	0.2	0.2	0.0	0.0
M	Scand?	42	1960.6	3.0	1.3830	0.0070	1962.5	1.9	1959.5	1959.3	-1.1	1.1	-1.3	1.3
M	TX	42	1960.1	3.0	1.5051	0.1504	1962.8	2.7	1959.8	1959.6	-0.3	0.3	-0.5	0.5
M	Scand?	42	1973.0	3.0	1.3951	0.0052	1974.7	1.7	1971.7	1971.5	-1.3	1.3	-1.5	1.5
M	Sweden	42	1989.1	3.0	1.1481	0.0043	1991.4	2.3	1988.4	1988.2	-0.7	0.7	-0.9	0.9
F	Sweden	42	1958.1	2.8	1.6568	0.0062	1966.8	8.7	1963.8	1963.6	5.7	5.7	5.5	5.5
M	Georgia	42	1981.1	3.0	1.2294	0.0046	1983.3	2.2	1980.3	1980.1	-0.8	0.8	-1.0	1.0
			Mean	3.0			Mean	3.2		Mean	0.2	1.4	0.0	1.5
							SD	2.5		SD	2.5	1.9	2.5	1.8
M	n/a	33	1962.6	4.3	1.5190	0.0060	1967.1	4.5	1962.8	1963.5	0.2	0.2	0.9	0.9
M	TX	43	1960.1	4.3	1.4475	0.0062	1962.8	2.7	1958.5	1959.2	-1.6	1.6	-0.9	0.9
M	Bosnia	33	1953.0	4.3	1.0126	0.0039	1956.1	3.1	1951.8	1952.5	-1.2	1.2	-0.5	0.5
M	Bosnia	43	1953.0	4.3	1.0634	0.0507	1957.2	4.2	1952.9	1953.6	-0.1	0.1	0.6	0.6
			Mean	4.3			Mean	3.6		Mean	-0.7	0.8	0.0	0.7
							SD	0.9		SD	0.9	0.7	0.9	0.2
M	Sweden	34	1991.8	5.1	1.1145	0.0036	1996.0	4.2	1991.1	1991.9	-0.7	0.7	0.1	0.1
M	Sweden	34	1988.6	5.1	1.1366	0.0076	1992.7	4.1	1987.8	1988.6	-0.8	0.8	0.0	0.0
M	Scand?	34	1990.0	5.1	1.1080	0.0040	1997.2	7.2	1992.3	1993.1	2.3	2.3	3.1	3.1
M	Sweden	34	1990.8	5.1	1.1212	0.0043	1995.0	4.2	1990.1	1990.9	-0.7	0.7	0.1	0.1
M	CT	34	1996.3	5.1	1.0872	0.0038	2001.0	4.7	1996.1	1996.9	-0.2	0.2	0.6	0.6
F	Sweden	34	1993.6	4.4	1.1008	0.0039	1998.4	4.8	1993.5	1994.3	-0.1	0.1	0.7	0.7
F	Sweden	34	1991.4	4.4	1.1196	0.0037	1995.2	3.8	1990.3	1991.1	-1.1	1.1	-0.3	0.3
F	TX	34	1991.4	4.4	1.1224	0.0032	1994.6	3.2	1989.7	1990.5	-1.7	1.7	-0.9	0.9
F	Sweden	44	1988.8	4.4	1.1381	0.0033	1992.2	3.4	1987.3	1988.1	-1.5	1.5	-0.7	0.7
F	Sweden	44	1992.5	4.4	1.1156	0.0042	1995.8	3.3	1990.9	1991.7	-1.6	1.6	-0.8	0.8

M	Sweden	44	1991.3	5.1	1.1218	0.0049	1995.0	3.7	1990.1	1990.9	-1.2	1.2	-0.4	0.4
M	Sweden	44	1993.2	5.1	1.1054	0.0035	1997.7	4.5	1992.8	1993.6	-0.4	0.4	0.4	0.4
M	BC	44	1996.6	5.1	1.1130	0.0042	1996.3	-0.3	1991.4	1992.2	-5.2	5.2	-4.4	4.4
M	CT	44	1997.2	5.1	1.0940	0.0039	2000.0	2.8	1995.1	1995.9	-2.1	2.1	-1.3	1.3
M	TX	44	1960.1	5.1	1.6666	0.0067	1966.4	6.3	1961.5	1962.3	1.4	1.4	2.2	2.2
M	Scand?	44	1962.6	5.1	1.5840	0.0070	1968.3	5.7	1963.4	1964.2	0.8	0.8	1.6	1.6
			Mean	4.9			Mean	4.1		Mean	-0.8	1.4	0.0	1.1
							SD	1.7		SD	1.7	1.2	1.7	1.2
F	Sweden	35	1989.2	5.7	1.1211	0.0037	1994.9	5.7	1988.9	1989.6	-0.3	0.3	0.4	0.4
M	Uruguay	35	1966.8	6.5	1.4720	0.0043	1972.4	5.6	1966.4	1967.1	-0.4	0.4	0.3	0.3
F	Scand?	45	1973.7	5.7	1.3231	0.0044	1978.4	4.7	1972.4	1973.1	-1.3	1.3	-0.6	0.6
F	Mexico	45	1963.8	5.7	1.5331	0.0058	1970.2	6.4	1964.2	1964.9	0.4	0.4	1.1	1.1
F	WA	45	1962.8	5.7	1.6356	0.0057	1967.4	4.6	1961.4	1962.1	-1.4	1.4	-0.7	0.7
M	Sweden	45	1992.2	6.5	1.1020	0.0039	1998.2	6.0	1992.2	1992.9	0.0	0.0	0.7	0.7
M	MA	45	1957.7	6.5	1.6698	0.0072	1962.0	4.3	1956.0	1956.7	-1.7	1.7	-1.0	1.0
			Mean	6.0			Mean	5.3		Mean	-0.7	0.8	0.0	0.7
							SD	0.8		SD	0.8	0.7	0.8	0.3
M	Chile	36	1967.8	2.4	1.5222	0.0118	1970.8	3.0	1968.4	1967.8	0.6	0.6	0.0	0.0
M	Scand?	36	1955.8	2.4	0.9934	0.0031	1955.1	-0.7	1952.7	1952.1	-3.1	3.1	-3.7	3.7
M	Mexico	36	1954.8	2.4	1.0738	0.0038	1957.4	2.6	1955.0	1954.4	0.2	0.2	-0.4	0.4
M	TX	36	1992.0	2.4	1.1019	0.0035	1998.2	6.2	1995.8	1995.2	3.8	3.8	3.2	3.2
F	Uruguay	36	1992.3	2.3	1.1158	0.0036	1995.7	3.4	1993.3	1992.7	1.0	1.0	0.4	0.4
F	Na	36	1963.5	2.3	1.6320	0.0070	1967.4	3.9	1965.0	1964.4	1.5	1.5	0.9	0.9
M	TX	46	1965.3	2.4	1.4075	0.0041	1962.6	-2.7	1960.2	1959.6	-5.1	5.1	-5.7	5.7
F	BC	36	1970.4	2.3	1.3631	0.0058	1976.6	6.2	1974.2	1973.6	3.8	3.8	3.2	3.2

F	Sweden	46	1976.6	2.3	1.2552	0.0054	1981.6	5.0	1979.2	1978.6	2.6	2.6	2.0	2.0
			Mean	2.4			Mean	3.0		Mean	0.6	2.4	0.0	2.2
							SD	3.0		SD	3.0	1.7	3.0	1.9
M	Scand?	37	1979.4	6.5	1.2200	0.0047	1983.9	4.5	1977.7	1977.7	-1.7	1.7	-1.7	1.7
M	Mexico	37	1954.8	6.5	1.6573	0.0059	1962.7	7.9	1956.5	1956.5	1.7	1.7	1.7	1.7
M	TX	37	1960.1	6.5	1.6735	0.0056	1966.8	6.7	1960.6	1960.6	0.5	0.5	0.5	0.5
M	WA	37	1970.6	6.5	1.3518	0.0048	1976.8	6.2	1970.6	1970.6	0.0	0.0	0.0	0.0
M	Scand?	37	1971.7	6.5	1.2710	0.0050	1980.8	9.1	1974.6	1974.6	2.9	2.9	2.9	2.9
F	Scand?	37	1963.7	5.6	1.4810	0.0060	1972.0	8.3	1965.8	1965.8	2.1	2.1	2.1	2.1
F	Sweden	47	1963.7	5.6	1.5644	0.0057	1969.6	5.9	1963.4	1963.4	-0.3	0.3	-0.3	0.3
F	Scand?	47	1977.8	5.6	1.2320	0.0050	1983.2	5.4	1977.0	1977.0	-0.8	0.8	-0.8	-0.8
M	Scand?	47	1962.4	6.5	1.6172	0.0060	1967.6	5.2	1961.4	1961.4	-1.0	1.0	-1.0	1.0
M	TX	47	1960.1	6.5	1.6945	0.0060	1966.5	6.4	1960.3	1960.3	0.2	0.2	0.2	0.2
M	WA	47	1973.4	6.5	1.3421	0.0047	1977.5	4.1	1971.3	1971.3	-2.1	2.1	-2.1	2.1
F	Irak	47	1967.6	5.6	1.4720	0.0055	1972.3	4.7	1966.1	1966.1	-1.5	1.5	-1.5	1.5
			Mean	6.2			Mean	6.2		Mean	0.0	1.2	0.0	1.1
							SD	1.6		SD	1.6	0.9	1.6	1.1
M	Marocko	38	1958.3	13.0	1.5362	0.0055	1970.0	11.7	1957.4	1959.0	-0.9	0.9	0.7	0.7
M	Scand?	38	1950.6	13.0	1.1802	0.0031	1957.8	7.2	1945.2	1946.8	-5.4	5.4	-3.8	3.8
M	WA	38	1982.3	13.0	1.1357	0.0040	1992.5	10.2	1979.9	1981.5	-2.4	2.4	-0.8	0.8
F	Scand?	38	1971.2	11.8	1.2452	0.0040	1982.4	11.2	1969.8	1971.4	-1.4	1.4	0.2	0.2
F	Chile	38	1955.9	11.8	1.5473	0.0054	1970.0	14.1	1957.4	1959.0	1.5	1.5	3.1	3.1
M	Quwait	48	1970.4	13.0	1.2715	0.0050	1981.0	10.6	1968.4	1970.0	-2.0	2.0	-0.4	0.4
M	Scand?	48	1957.4	13.0	1.5586	0.0063	1969.3	11.9	1956.7	1958.3	-0.7	0.7	0.9	0.9
M	Scand?	48	1964.9	13.0	1.3236	0.0040	1978.3	13.4	1965.7	1967.3	0.8	0.8	2.4	2.4

M	WA	48	1982.3	13.0	1.1323	0.0042	1993.1	10.8	1980.5	1982.1	-1.8	1.8	-0.2	0.2
F	Scand?	48	1961.7	11.8	1.5186	0.0063	1970.9	9.2	1958.3	1959.9	-3.4	3.4	-1.8	1.8
			Mean	12.6			Mean	11.0		Mean	-0.8	1.5	0.0	1.1
							SD	1.9		SD	1.7	1.1	1.5	1.0

¹In the four studies, teeth were collected by dentists in different countries. Information about country of origin was not recorded by Swedish dentists in the first publication (6), but n/a typically implies that the subject either was raised in Europe or Middle East. Scand? = probably raised in Scandinavia.

²Enamel formation time according to Nolla (25).

³DOB of the person was calculated by subtracting the actual DOB from the average estimated C-14 incorporation time. When doing so, the average error mathematically becomes 0.0 in the second last column.

⁴Unclear when this subject moved to WA.