LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

LLNL-PROC-669626

TRIPOLI-4 and FREYA for Stochastic Analog Neutron Transport. Application to Neutron Multiplicity Counting.

J. M. Verbeke, O. Petit

April 15, 2015

TRIPOLI-4 and FREYA for Stochastic Analog Neutron Transport. Application to Neutron Multiplicity Counting.
Manchester, United Kingdom
May 19, 2015 through May 21, 2015

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

TRIPOLI-4® and FREYA for Stochastic Analog Neutron Transport. Application to Neutron Multiplicity Counting.

J. M. Verbeke*
Lawrence Livermore National Laboratory
P.O. Box 808, Livermore, California 94551, USA

O. Petit
Commissariat à l'Energie Atomique et aux Energies Alternatives
CEA-Saclay, DEN, DM2S, SERMA, LTSD
F-91191 Gif-sur-Yvette Cedex, France

Abstract

From nuclear safeguards to homeland security applications, the need for better modeling of nuclear interactions has grown over the past decades. Current Monte Carlo radiation transport codes compute average quantities with great accuracy and performance, but performance and averaging come at the price of limited interaction-by-interaction modeling. These codes often lack the capability of modeling interactions exactly: for a given collision, energy is not conserved, energies of emitted particles are uncorrelated, multiplicities of prompt fission neutrons and photons are uncorrelated. Many modern applications require more exclusive quantities than averages, such as the fluctuations in certain observables (e.g. the neutron multiplicity) and correlations between neutrons and photons. In an effort to meet this need, the radiation transport Monte Carlo code TRIPOLI-4® was modified to provide a specific mode modeling nuclear interactions in a full analog way, replicating as much as possible the underlying physical process. Furthermore, the computational model FREYA (Fission Reaction Event Yield Algorithm) was coupled with TRIPOLI-4® to model complete fission events. FREYA automatically includes fluctuations as well as correlations resulting from conservation of energy and momentum.

Neutron Multiplicity Counting (NMC) exploits the correlated nature of fission chains, and thus requires analog neutron transport. With the latest analog neutron transport developments in TRIPOLI-4®, we will show that NMC can now be properly simulated, by reconstructing the mass and multiplication of an object by analyzing the measured signal from ^{3}He tubes in a well counter.

1 INTRODUCTION

Methods based on time-correlated signals have been developed over many years to characterize fissile materials. For NMC, sequences of thermal neutron captures are recorded in ^{3}He tubes. To determine features of the measured objects, the sequences are split into time windows, and the numbers of neutrons arriving in each window are recorded to build statistical count distributions. These distributions are in turn analyzed to authenticate or characterize fissile materials. Some materials such as ^{252}Cf emit several neutrons simultaneously, whereas others such as uranium and plutonium multiply the number of neutrons to form bursts. This translates into unmistakable time-correlated signatures.

*This report was written while J.M. Verbeke was on a professional leave at CEA-Saclay.

General Monte Carlo codes that are used for criticality safety evaluations are typically meant for calculation of an integral reactor parameter such as k_{eff} and for estimation of neutron fluxes and derived quantities of interest. They make use of well established variation reduction techniques leading to more efficient calculations. These techniques are meant to speed up calculations and are sufficient for the calculation of average quantities such as flux, energy deposition and multiplication. However, they suffer from approximations of the underlying physical interactions, and are thus unsuitable for studying detailed correlations between neutrons and/or photons on an interaction-by-interaction basis, and in particular for NMC which relies on the correlated nature of fission chains.

The first part of this paper will focus on the latest TRIPOLI-4®[1,2] developments that were necessary to simulate NMC experiments: analog neutron transport, coupling with the LLNL Fission Library/FREYA[3–5] package for fission interactions, development of a spontaneous fission source, and new options to reduce memory footprint of ROOT[6] track files. The second part will focus on the use of these new capabilities for NMC. We will show that the mass and multiplication of a PuO_2 ball in a well counter (see Fig. 1) can be determined from measurements of the neutron captures in the ^{3}He tubes.



(a) View 1, multiple ^{3}He tubes removed for clarity. (b) View 2, upper polyethylene plug removed.

Figure 1: (color online) Cutouts to show the inside of a well counter. Polyethylene (magenta), Cadmium (cyan), ^{3}He tubes (green), representation of a generic neutron source to be characterized (blue).

2 DEVELOPMENTS IN TRIPOLI-4® FOR NMC

TRIPOLI-4® solves the linear Boltzmann equation for neutrons, photons, electrons and positrons with the Monte Carlo method, in any 3-D geometry. The code uses ENDF format continuous-energy cross sections from various international evaluations. It has advanced variance reduction methods to address deep penetration issues and can be run in parallel. TRIPOLI-4® is used as a reference code for industrial purposes (fission/fusion) for CEA[a], EDF[b] and branches of AREVA, as well as an R&D and teaching tool, for radiation protection and

[a]Commissariat à l'énergy atomique et aux énergies alternatives.
[b]Electricité de France.

shielding, core physics, nuclear criticality safety and nuclear instrumentation.

This section presents the list of the most important TRIPOLI-4® developments that were required for the NMC application, starting from version 9 of the code. These developments were made using a recent version of TRIPOLI-4® with analog mode capabilities.[7] While simulating the well counter was the objective of this study, these developments would also apply to multiplicity counting with liquid scintillators for fast neutrons and photons.

2.1 Coupling of FREYA and TRIPOLI-4® for fission modeling

To model fission, general-purpose Monte Carlo codes (TRIPOLI-4®, MCNP6/X,[8] TART,[9] COG,[10] Geant,[11] etc.) employ the "average fission model" which is characterized by outgoing projectiles (fission neutrons and photons) that are uncorrelated, and sampled from the same probability density functions.

During the past decade several code extensions have been developed that allow the modeling of correlations in fission. MCNP-DSP[12] and MCNPX-PoliMi[13] added limited angular correlations of fission neutrons. The LLNL Fission Library,[14] introduced in MCNPX2.7.0,[15] Geant 4.9[11] and MCNP6, featured time-correlated sampling of photons from neutron-induced fission, photofission and spontaneous fission. The capabilities for correlations are, however, limited as they sample outgoing particles from average fission distributions instead of sampling them from individual realizations of a fission process.

In recent years, various simulation treatments addressed fluctuations of and correlations between fission observables. In particular, a Monte Carlo approach was developed[16] for the sequential emission of neutrons and photons from individual fission fragments in binary fission. The more recent event-by-event fission model, FREYA, has been specifically designed for producing large numbers of fission events in a fast simulation.[17] Employing nuclear data for fragment mass and kinetic energy distributions, using statistical evaporation models for neutron and photon emission, and conserving energy, momentum, and angular momentum throughout, FREYA is able to predict a host of correlation observables, including correlations in neutron multiplicity, energy, and angles, and the energy sharing between neutrons and photons. For modeling of fission on an interaction-by-interaction basis, the new LLNL Fission Library/FREYA package was coupled with TRIPOLI-4®.

2.2 Development of a spontaneous fission source

A spontaneous fission source was developed to sample time-correlated neutrons and photons from fission. This source emits bursts of time-correlated prompt neutrons and photons from individual fission events, whose multiplicities and energies are sampled from the LLNL Fission Library/FREYA package. TRIPOLI-4® accesses this source as an *external source* (see User Manual[1]). The times of spontaneous fissions are sampled randomly and uniformly within a given time interval ΔT[c]. The rate of spontaneous fissions has to match the rate F_s of spontaneous fissions of the object to be measured experimentally. It is therefore essential to set the correct number of particles accordingly.

2.3 Reduction in tracks memory footprint

To model NMC for the well counter shown in Fig. 1, it is necessary to store the time tags of all the neutron capture reactions in the ^{3}He tubes.

It was quickly realized that the ROOT tracks stored by TRIPOLI-4® became bloated for large simulations, leading to files that were close to terabytes in size for seconds of experimental data. Most of the tracks did not result in ^{3}He(n,p) reactions and were thus cluttering the disk. When filtering out tracks failing to traverse detector cells, we were able to substantially decrease the memory footprint, but not enough. Two new options

[c] The time tags of the ^{3}He(n,p) reactions in the ^{3}He tubes are re-ordered chronologically in post-processing.

were therefore introduced to further reduce the size of the track files. The first option enables us to store full tracks containing one or several events of interest, whereas the second option enables storage of only specific events with those tracks. With these two additional filters, we could keep the footprint of the ROOT track files in check.

3 PUO_2 BALL

Let's consider a PuO_2 object spherical in shape, of weight equal to 5.5366 kg, density 3 g/cc, and of outer radius 7.62 cm[d]. Knowing the neutron yields of the different isotopes composing the object, we calculated the spontaneous fission rate to be 140,170 spontaneous fissions/s. The spontaneous fission source is uniformly distributed across the sphere. A simulation of the PuO_2 source in the well counter shown in Fig. 1 was performed. While the intensity of the (α,n) source could be calculated, we will neglect this contribution for the purpose of this study.

3.1 Fitting count distributions to determine system parameters

The arrival times of the neutrons in each of the ^{3}He tubes were recorded in the simulation. Randomly splitting the sequence of time tags into N segments of width T — where T is of the order of microseconds to hundreds of microseconds — one can count how many neutrons arrive in the first segment, how many in the second segment, in the third one, etc. to build a distribution $B_n(T)$ of the number n of neutrons arriving in the segments of width T. The blue dots labeled "simulated data" in Fig. 4(a) show a typical count distribution.

The probability distributions $b_n(T)$ ($b_n(T)$ is the probability of recording n counts in a time gate T, which is equivalent to $B_n(T)$ normalized by the number of segments of width T) can be reconstructed theoretically[18] using different sets of the 3 free parameters (M, ε, $\bar{\nu}_{\rm sp}F_s$), where M is the multiplication of the object, ε the efficiency of the detector array, $\bar{\nu}_{\rm sp}$ the average number of neutrons emitted per spontaneous fission and F_s the intensity in units of spontaneous fissions per second of the spontaneous fission sources in the object. Using a likelihood function, one can determine which parameters (M, ε, $\bar{\nu}_{\rm sp}F_s$) generate the theoretical count distribution $b_n^{\rm theory}(T)$ closest to the measured data points $b_n(T)$. This method is best described in Ref.[19]

Each set of parameters (M, ε, $\bar{\nu}_{\rm sp}F_s$) has an associated likelihood that the reconstructed $b_n^{\rm theory}(T)$ will be a good match to the measured $b_n(T)$. Using Bayes' theorem, we calculate the posterior probability of each such set. To determine the region of the (M, ε, $\bar{\nu}_{\rm sp}F_s$) space that contains the solution with a credibility of 68.27%, we have to accumulate high posterior probability sets until the cumulative probability reaches 68.27%. Fig. 2 shows the credible regions in the (M, ε) and (M, $\bar{\nu}_{\rm sp}F_s$) parameter spaces for credibilities of 68.27% (red), 95.45% (yellow) and 99.73% (blue). The top 2 graphs are computed with FREYA. The bottom 2 graphs are computed without FREYA. Without FREYA, $\bar{\nu}$ is statistically rounded up or down at each fission site to get a number of neutrons. Fig. 3 shows the same credible regions for 2530 seconds.

[d]The isotopics of the plutonium are 0.014% ^{238}Pu, 93.5% ^{239}Pu, 6% ^{240}Pu, 0.5% ^{241}Pu, 0.03% ^{242}Pu, and traces of other isotopes



(a) (M, ε) with FREYA.

(b) (M, $\bar{\nu}_{\mathrm{sp}}F_s$) with FREYA.

(c) (M, ε), without FREYA.

(d) (M, $\bar{\nu}_{\mathrm{sp}}F_s$), without FREYA.

Figure 2: (color online) Credible regions for theoretical reconstructions: 68.27% (red), 95.45% (yellow), 99.73% (blue). Nuclear data for induced fission of ^{239}Pu at 1 MeV. The measurement time is equivalent to 350 seconds.



(a) (M, ε) with FREYA.

(b) (M, $\bar{\nu}_{\mathrm{sp}} F_s$) with FREYA.

(c) (M, ε) without FREYA.

(d) (M, $\bar{\nu}_{\mathrm{sp}} F_s$) without FREYA.

Figure 3: (color online) Credible regions for theoretical reconstructions of PuO_2 ball: 68.27% (red), 95.45% (yellow), 99.73% (blue). Nuclear data for induced fission of ^{239}Pu at 1 MeV. The measurement time is equivalent to 2530 seconds.

3.2 Discussions

Table 1 shows the multiplication calculated by TRIPOLI-4® in different modes: either a criticality calculation including a convergence process of the fission source or a fixed source criticality calculation where the same neutron source is kept during the whole simulation and fission neutrons are sampled but not used for the source convergence (see TRIPOLI-4® User Manual[1]).

Table 1: Neutron multiplication for PuO_2 ball in well counter calculated with TRIPOLI-4®.

mode	LLNL Fission Library/FREYA	M$\pm\sigma$
CRITICALITY	no	1.3312 ± 0.0012[e]
FIXED_SOURCES_CRITICALITY	no	1.3140 ± 0.0007
FIXED_SOURCES_CRITICALITY	yes	1.3124 ± 0.0008

With TRIPOLI-4® running in analog mode, the multiplication of the PuO_2 ball within the well counter was calculated to be 1.3124 ± 0.0008 with FREYA, and 1.3140 ± 0.0007 without FREYA. These multiplications are very close and show that average quantities are not affected by the choice of the fission model, whether it statistically samples $\bar{\nu}$ rounded up or down, or a full neutron multiplicity distribution.

For the reconstruction with FREYA, the best solution is (M, ε, $\bar{\nu}_{sp}F_s$)=(1.35, 20.9%, 308368 n/s). One may wonder whether the credible regions shown in Fig. 3 contain the true solution, which is the one with the source intensity used for the simulation and the multiplication computed by TRIPOLI-4®. The solution (1.33, 21.5%, 303041 n/s) in Figs. 3(a)-3(b) is within the 68.27% credible region and gives the correct source intensity of 302799 n/s. The small discrepancies are likely to be attributed to systematic errors, and to inadequacies in the theory to model the experiment. This is discussed at length in Ref.[19]

It is interesting to compare the count distributions reconstructed from the different solutions within the 68.27% credible region. A set of 3 such distributions is shown in Fig. 4(a). All the solutions within the credible region are essentially indistinguishable, which explains the size of the uncertainty in that region, and illustrates the highly degenerate nature of the model.

Without FREYA, the best solution for the reconstruction is (M, ε, $\bar{\nu}_{sp}F_s$)=(1.16, 27.9%, 257174 n/s). Within the 68.27% credible region, the solution that gives the closest source intensity is (M, ε, $\bar{\nu}_{sp}F_s$)=(1.2, 26.0%, 269471 n/s). The model still gives very good count distribution reconstructions, as Fig. 4(b) illustrates, but unfortunately, these reconstructions are for the incorrect parameters. For correlated quantities and low multiplication, sampling the full distribution for the fission neutron multiplicity is important. The correct solution could be found with FREYA, whereas without FREYA the default neutron multiplicity sampling gave incorrect solutions.

4 CONCLUSION

FREYA was coupled to TRIPOLI-4® for the purpose of using the latter for NMC. With the addition of a few modifications such as analog transport, spontaneous fission sources and improved tracking capabilities, we

[e] This multiplication was calculated using $M = 1/(1-k_{\text{eff}})$ and the k-eigenvalue method of TRIPOLI-4® for k_{eff}. Since this method does not solve the same problem as the FIXED_SOURCES_CRITICALITY method,[20] it is not expected to produce the same multiplication. It is only shown for the sake of completeness.



(a) with FREYA (b) without FREYA

Figure 4: (color online) Comparison between theoretically reconstructed count distributions within the 68.27% credible region in Fig. 3. PuO_2 ball. Random time gate count distribution. Time gate width=1 ms. Simulation result in blue. Nuclear data for induced fission of ^{239}Pu at 1 MeV. The measurement time is equivalent to 2530 seconds.

demonstrated by way of a PuO_2 ball simulation that TRIPOLI-4® when coupled with FREYA can simulate physical correlations sufficiently well to reproduce predicted count distributions measured by a well counter.

Average quantities like neutron flux, reaction rate, multiplication are not affected by the choice of the fission model. Whether the fission model statistically samples $\bar{\nu}$ rounded up or down, or a full fission neutron multiplicity distribution, to emit a number of secondary neutrons, has little to no impact on the result.

For methods using correlated quantities, sampling the full distribution for the fission neutron multiplicity is paramount. The correct solution to the PuO_2 ball problem could be found with the fission model FREYA, whereas the default neutron multiplicity sampling gave incorrect solutions[f]. Thus including such capabilities in Monte Carlo transport codes is important.

ACKNOWLEDGMENTS

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

TRIPOLI-4® is a registered trademark of CEA. We gratefully acknowledge EDF for their long term partnership and AREVA for their support.

J.M. Verbeke wishes to acknowledge the Office of Defense Nuclear Nonproliferation Research and Development in DOE/NNSA for their support, and to thank the stochastic team (LTSD) of the Service d'Etudes de Réacteurs et de Mathématiques Appliquées (SERMA) in the Commissariat à l'Energie Atomique et aux Energies Alternatives at Saclay, France for hosting him while this paper was being written and while FREYA was being coupled with TRIPOLI-4®.

[f] We could show that the choice of the fission multiplicity distribution becomes less important the higher the multiplication, but the theoretical developments required for this are beyond the scope of this paper.

References

[1] TRIPOLI-4® Project Team: "TRIPOLI-4 version 8 User Guide," CEA-R-6316 (2013), see also http://www.nea.fr/abs/html/nea-1716.html. 2, 3, 7

[2] TRIPOLI-4® Project Team: "TRIPOLI-4, CEA, EDF and AREVA Reference Monte Carlo Code," Joint Int. Conf. on SNA and MC 2013 (SNA+MC 2013), Paris, France (2013), also accepted for publication in Annals of Nuclear Energy, 2015. 2

[3] J.M. VERBEKE, J. RANDRUP, R. VOGT, "Fission Reaction Event Yield Algorithm FREYA for Event-by-Event Simulation of Fission," to be published in Computer Physics Communications. 2

[4] C. HAGMANN, J. RANDRUP, R. VOGT, "FREYA - A new Monte Carlo code for improved modeling of fission chains," Trans. Nucl. Sci. **60**, 545-549 (2013). 2

[5] J. M. VERBEKE, C.A. HAGMANN, J. RANDRUP, R. VOGT, "Integration of FREYA into MCNP6: An Improved Fission Chain Modeling Capability," Lawrence Livermore National Laboratory, Livermore, CA, LLNL-PROC-638986 (2013). 2

[6] R. BRUN, F. RADEMAKERS, "ROOT - An Object Oriented Data Analysis Framework," Proceedings AIHENP'96 Workshop, Lausanne, Sep. 1996, Nucl. Inst. Meth. Phys. Res. A **389**, 81-86 (1997). See also http://root.cern.ch/. 2

[7] O. PETIT, E. DUMONTEIL, "Analog neutron transport for nuclear instrumentation applications with the Monte-Carlo code TRIPOLI-4," ANS RPSD 2014 - 18th Topical Meeting of the Radiation Protection & Shielding Division of ANS Knoxville, TN, September 14 - 18, 2014, American Nuclear Society, LaGrange Park, IL (2014). 3

[8] T. GOORLEY *et al.* "Initial MCNP6 Release Overview - MCNP6 version 1.0," Los Alamos National Laboratory, Los Alamos, NM, LA-UR-13-22934 (2013). 3

[9] D. E. CULLEN, "TART2005: A Coupled Neutron-Photon 3-D, Combinatorial Geometry, Time Dependent Monte Carlo Transport Code," Lawrence Livermore National Laboratory, Livermore, CA, UCRL-SM-218009 (2005). 3

[10] R. M. BUCK, E. M. LENT, "COG User's Manual: A Multiparticle Monte Carlo Transport Code," Lawrence Livermore National Laboratory, Livermore, CA, UCRL-TM-202590, 5th Edition (2002). 3

[11] Available from the European Organization for Nuclear Research, http://geant4.cern.ch. 3

[12] T. E. VALENTINE, "MCNP-DSP Users Manual," Oak Ridge National Laboratory, Oak Ridge, TN, ORNL/TM-13334 R2 (2001). 3

[13] E. PADOVANI, S. A. POZZI, S. D. CLARKE, E. C. MILLER, "MCNPX-PoliMi User's Manual," C00791 MNYCP, Radiation Safety Information Computational Center, Oak Ridge National Laboratory (2012). 3

[14] J. M. VERBEKE, C. HAGMANN, D. WRIGHT, "Simulation of Neutron and Gamma-Ray Emission from Fission and Photofission," Lawrence Livermore National Laboratory, Livermore, CA, UCRL-AR-228518 (2010). 3

[15] D. B. PELOWITZ *et al.*, "`MCNPX2.7.0` Extensions," Los Alamos National Laboratory, Los Alamos, NM, LA-UR-11-02295 (2011). 3

[16] S. LEMAIRE, P. TALOU, T. KAWANO, M. B. CHADWICK, D. G. MADLAND, "Monte Carlo approach to sequential γ-ray emission from fission fragments," Phys. Rev. C **73**, 014602 (2006). 3

[17] R. VOGT, J. RANDRUP, "Event-by-event study of photon observables in spontaneous and thermal fission," Phys. Rev. C **87**, 044602 (2013). 3

[18] M. K. PRASAD, N. J. SNYDERMAN, "Statistical Theory of Fission Chains and Generalized Poisson Neutron Counting Distributions," Nucl. Sci. Eng. **172**, 300 (2012). 4

[19] J. M. VERBEKE, "Neutron Multiplicity Counting: Confidence Intervals for Reconstruction Parameters," Lawrence Livermore National Laboratory, Livermore, CA, LLNL-JRNL-668401-DRAFT, submitted to Nuclear Science and Engineering (2015). 4, 7

[20] D.E. CULLEN, C.J. CLOUSE, R. PROCASSINI, R.C. LITTLE, "Static and Dynamic Criticality: Are They Different," Lawrence Livermore National Laboratory, Livermore, CA, UCRL-TR-201506 (2003). 7