



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

On the v -Representability in Density Functional Theory: Application to Non-interacting Systems

M. Daene, A. Gonis

June 8, 2016

Computation

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

On the v -Representability Problem in Density Functional Theory: Application to Non-interacting Systems

M. Däne^{1,†,‡} and A. Gonis^{1,‡}

¹ Lawrence Livermore National Laboratory; daene1@llnl.gov

² Lawrence Livermore National Laboratory; gonis1@llnl.gov

* Correspondence: daene1@llnl.gov; Tel.: +1-925-423-8181

† Current address: Affiliation 1 and 2: Lawrence Livermore National Laboratory, PO Box 808, L-372, Livermore, CA 94551

‡ These authors contributed equally to this work.

Academic Editor: Prof. Karlheinz Schwarz and Prof. Agnes Nagy

Abstract: Based on a computational procedure for determining the functional derivative with respect to the density of *any* antisymmetric N -particle wave function for a non-interacting system that leads to the density, we devise a test as to whether or not a wave function known to lead to a given density corresponds to a solution of a Schrödinger equation for some potential. We examine explicitly the case of non-interacting systems describe by Slater determinants. Numerical examples for the cases of a one-dimensional square-well potential with infinite walls and the harmonic oscillator potential illustrate the formalism.

Keywords: density functional theory; v -representability; constrained search

1. Introduction

The Hohenberg-Kohn theorems [1] form the foundation of density functional theory (DFT), on which most *ab-initio* electronic structure methods currently are based. The first theorem proves that the ground state of an many-electron system, described by an external potential, is uniquely determined by the ground state density (within an arbitrary constant). Therefore, the ground state density is the independent variable of DFT. In practice, it is still challenging to find the many-body wave function for the ground state of a potential. That task has been simplified by Kohn and Sham [2] who introduced a non-interacting system (Kohn-Sham system) with the same density as the interacting system that minimized the kinetic energy only. The existence of such a system is only assumed, but essential to the Kohn-Sham formulation of DFT.

In most of today's application of DFT only one direction of the Hohenberg-Kohn theorems is used, to find the ground state density for a given system described by an external potential. For the other direction, the inverse problem, it is a-priori not clear if even a potential exists that leads to a given density through the solution of the Schrödinger equation, whether for the interacting or non-interacting system, the problem of v -representability. We refer to the literature [3–11] for discussions on the issue of v -representability. Specific examples of v -representable densities as well as non- v -representable densities have being identified.

An alternative formulation of the problem can be developed based on the N -representability property [3–6] of any density normalized to an integral number of particles, namely that a density can always be obtained from a wave function for a state of N -particles that, for fermions, is antisymmetric with respect to interchange of individual particle coordinates (and spins for electrons). The v -representability problem can be stated as follows: Given a density corresponding to a minimizing wave function, as obtained through the constrained search [3], determine whether or not the wave function is derived as the solution of the Schrödinger equation for a system of N electrons, either interacting, or non-interacting. A more general question is concerned with whether or not a

34 wave function that leads to a given density is either interacting or non-interacting v -representable.
 35 A formal procedure that addresses this problem for general interacting systems has been published
 36 elsewhere [12]. In the following, a density defined with the following properties: it is everywhere
 37 non-negative, is normalized to the number of particles, and leads to a finite value of the kinetic energy.

38 In this work we focus on the v -representability for non-interacting systems, even though all
 39 procedures presented here can be applied for the interacting system in the same way [12]. In previous
 40 work [13–15], we have derived closed expressions for the functional derivative of terms that only
 41 depend indirectly on the density, with respect to it. Using that formalism, we now show analytically
 42 that differentiating the non-interacting kinetic energy for a v -representable wave function leads to the
 43 negative potential (Kohn-Sham potential), up to a constant. This does not hold for arbitrary wave
 44 functions leading to the density. We obtain different wave functions leading to the same density
 45 by using a procedure by Cioslowski [16,17], who also showed that each of these wave functions is
 46 differentiable with respect to the density. This provides a test establishing whether a given wave
 47 function (non-interacting), leading to a density, is non-interacting v -representable. We illustrate these
 48 procedures on the one-dimensional particle in a box problem as well as the harmonic oscillator.

49 2. Results

50 2.1. Background

The Hamiltonian describing an interacting system of N electrons in an external potential takes the usual form,

$$\hat{H}^N = \hat{V} + \hat{T}^N + \hat{U}^N, \quad (1)$$

with the operators \hat{V} , \hat{T}^N and \hat{U}^N corresponding, respectively, to the external field, the kinetic energy and the inter-particle interaction (Coulomb repulsion for electrons). The ground-state energy of the system is given by the expectation value,

$$E_g = \langle \Psi_g^N | \hat{H}^N | \Psi_g^N \rangle, \quad \langle \Psi_g^N | \Psi_g^N \rangle = 1, \quad (2)$$

where $|\Psi_g^N\rangle$ denotes the many-particle ground state of \hat{H}^N . We use the notation, $|\Psi^N\rangle \rightarrow n(\mathbf{r})$, and say $|\Psi^N\rangle$ leads to the density $n(\mathbf{r})$, to denote the property,

$$n(\mathbf{r}) = N \int |\Psi^N(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_N)|^2 d\mathbf{r}_2 \dots d\mathbf{r}_N, \quad (3)$$

where $n(\mathbf{r})$ denotes the single-particle density function normalized to the total number of particles, N . We now write Eq. (2) in the form,

$$E_g = \int v(\mathbf{r}) n_g(\mathbf{r}) d\mathbf{r} + \langle \Psi_g^N | \hat{T} + \hat{U} | \Psi_g^N \rangle = \underbrace{\text{Min}}_{n(\mathbf{r})} \left[\int v(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} + F[n] \right] = \underbrace{\text{Min}}_{n(\mathbf{r})} E[n], \quad (4)$$

in terms of the constrained search functional [3,18],

$$F[n] = \underbrace{\text{Min}}_{|\Psi\rangle \rightarrow n(\mathbf{r})} \langle \Psi | \hat{T}^N + \hat{U}^N | \Psi \rangle. \quad (5)$$

51 Given a density, $n(\mathbf{r})$, the constrained search examines all antisymmetric N -particle wave functions
 52 that lead to the density and delivers the state (in the absence of degeneracy) that produces the
 53 minimum value of $\langle \Psi^N | \hat{T}^N + \hat{U}^N | \Psi^N \rangle$.

As in the initial formulation of DFT by Kohn and Sham [2], we postulate the existence of a fictitious non-interacting N -particle system described by the Hamiltonian,

$$\hat{H}_s^N = \hat{V}_s + \hat{T}^N, \quad (6)$$

under the action of an external potential, \hat{V}_s , whose ground-state density is identical to the density of the interacting system described by \hat{H}^N . In analogy with Eq. (5), we define the constrained search functional [3],

$$\begin{aligned} T_s[n] &= \underset{|\Phi^N\rangle \rightarrow n(\mathbf{r})}{\text{Min}} \langle \Phi^N | \hat{T}^N | \Phi^N \rangle = \langle \Phi_{\text{GS}}^N | T^N | \Phi_{\text{GS}}^N \rangle \\ &= \int \Phi^{N*}(\mathbf{r}_{(N)}) \left| \sum_j \left[-\frac{1}{2} \nabla_{\mathbf{r}_j}^2 \right] \right| \Phi^N(\mathbf{r}_{(N)}) \, d\mathbf{r}_{(N)}. \end{aligned} \quad (7)$$

54 implying a search over all Slater determinants leading to a given density and returning that
55 determinant that minimizes the expectation value of the kinetic energy for a system of N particles.

Now, assume that the density is non-interacting v -representable for some potential, $v_s(\mathbf{r})$, so that from the second theorem of Hohenberg and Kohn [1] follows that,

$$\frac{\delta T_s[n]}{\delta n(\mathbf{r})} = -v_s(\mathbf{r}), \quad (8)$$

defined within an arbitrary constant. The potential, $v_s(\mathbf{r})$, is such that the orbitals entering the construction of $\Phi^N(\mathbf{r}_{(N)})$ arise through the solutions of a single-particle Schrödinger equation,

$$\left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 + v_s(\mathbf{r}) \right] \phi_j(\mathbf{r}) = \epsilon_j \phi_j(\mathbf{r}), \quad (9)$$

with eigenenergies ϵ_j . Clearly, the orbitals minimize the expectation value of the kinetic energy and as such form the Slater determinant identified by the constrained search in establishing $T_s[n]$. For the case of Slater determinants, Eq. (7) reduces to the direct sum,

$$T_s[n] = \sum_j \int \phi_j^*(\mathbf{r}) \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \right] \phi_j(\mathbf{r}) \, d\mathbf{r}. \quad (10)$$

56 2.2. Functional Differentiation

57 The process of performing the functional differentiation of expectation values with respect to
58 the density, however, is not immediately evident: The dependence of a given wave function on the
59 density is implicit, rather than explicitly displayed in terms of the density (or analytic functions of
60 it). This difficulty is resolved through the use of the *equidensity orbitals* [19–21] that are written as
61 explicit functionals of the density and, as shown in [21], form an orthonormal and complete basis
62 in three-dimensional space. This property has been exploited in previous work [13–15] to allow the
63 functional differentiation of the Coulomb energy associated with a Slater determinant with respect to
64 the density in order to determine the Coulomb potential including the contribution of the exchange
65 term. A complete discussion is given in [14]. Here, we apply that formalism to derivatives of the
66 non-interacting kinetic energy with respect to the density.

We can obtain the functional derivative of $T_s[n]$ by differentiating the wave functions under the integral signs in (7), and determine the potential-like quantity,

$$-v_s(\mathbf{r}) = \frac{\delta T_s[n]}{\delta n(\mathbf{r})} = \left\langle \frac{\delta \Phi^N}{\delta n(\mathbf{r})} \left| \hat{T}^N \right| \Phi^N \right\rangle + \left\langle \Phi^N \left| \hat{T}^N \right| \frac{\delta \Phi^N}{\delta n(\mathbf{r})} \right\rangle, \quad (11)$$

67 where the double bars prohibit the operators from acting on the quantity beyond them. The reason
 68 for this is that functional derivatives of wave functions with respect to the density are generally not
 69 elements of the Hilbert space. Because of this feature, the operators in the last two equations act on
 70 the wave functions, either to their left or their right. It is seen that the combination of the constrained
 71 search and functional differentiation allows in principle the determination of all Slater determinants
 72 associated with given density that may correspond to a potential.

For the case of a Slater determinant, the last expression takes the form,

$$\frac{\delta T_s[n]}{\delta n(\mathbf{r})} = \sum_j \int \frac{\delta \phi^*(\mathbf{r}')}{\delta n(\mathbf{r})} \left[-\frac{1}{2} \nabla_{\mathbf{r}'}^2 \right] \phi_j(\mathbf{r}') \, d\mathbf{r}' + \sum_j \int \frac{\delta \phi(\mathbf{r}')}{\delta n(\mathbf{r})} \left[-\frac{1}{2} \nabla_{\mathbf{r}'}^2 \right] \phi_j^*(\mathbf{r}') \, d\mathbf{r}' = -v_s(\mathbf{r}), \quad (12)$$

73 where the del operator acts on arguments to its right, and the last line gives the functional derivative
 74 as a function of the coordinates that may or may not correspond to a potential. In this expression, the
 75 quantity, $T_s[n]$, denotes the expectation value of the non-interacting kinetic energy operator, \hat{T}^N , with
 76 respect to any wave function that leads to a density.

In previous work [13–15], we derived a closed-form expression of the functional derivative of
 an orbital $f(\mathbf{r})$ with respect to the density, $\frac{\delta f(\mathbf{r})}{\delta n(\mathbf{r})}$, in terms of ordinary (spatial) derivatives. In the
 one-dimensional case, excluding terms that would lead to a constant shift in the potential (constants
 and expressions solely depending on x but not x'), the derivative reads

$$\frac{\delta f(x)}{\delta n(x')} = \frac{f(x)}{2n(x)} \delta(x - x') + \frac{\Theta(x - x')}{n(x)} \left[f'(x) - \frac{f(x) n'(x)}{2n(x)} \right]. \quad (13)$$

77 Primes on functions denote spatial derivatives with respect to x and Θ is the Heaviside step function.
 78 Expressions for the three-dimensional case can be found in reference [14] and [15] for both the cases
 79 of cartesian and spherical coordinates, respectively.

80 In Appendix A we derive an expression for the differentiation of the expression for the kinetic
 81 energy (one-dimensional for simplicity) using equations (12) and (13), see Equation (25). If all orbitals
 82 contributing to the kinetic energy originate from the same potential through the Schrödinger or
 83 Kohn-Sham equations, we show analytically that the differentiation leads to the negative potential
 84 (within an arbitrary constant). The three-dimensional case follows analogously.

85 2.3. Illustration of the Formalism

86 The calculations quoted here illustrate the formalism developed above for cases where the
 87 potential is given and the wave functions are known analytically. The goal is to reproduce the
 88 potential from the functional derivative of the kinetic energy with respect to the density.

89 For simplicity, we discuss one-dimensional systems in the non-interacting case where the wave
 90 function can be represented as a Slater determinant. Analogous procedures will apply in higher
 91 dimensions and the interacting framework.

For our first example we consider N non-interacting particles in a box of length L bounded
 by potential walls of infinite height. The wave functions of the particles confined in the box are
 determined through the single-particle Schrödinger equation,

$$\left[-\frac{1}{2} \frac{d^2}{dx^2} + v(x) \right] \phi_n(x) = \epsilon_n \phi_n(x), \quad (14)$$

and vanish at the edges of the box. The ϵ_n denote the eigenvalues. When $v(x) = 0$ (or a constant), the
 normalized single-particle wave functions are given by the expressions,

$$\phi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) = A_n \sin(k_n x) \quad 0 \leq x \leq L \quad (15)$$

with quantum numbers $n = 1, 2, 3, \dots$. The energies take the form,

$$\epsilon_n = \frac{\hbar^2}{2m} \left(\frac{n\pi}{L} \right)^2 = \frac{\hbar^2}{2m} k_n^2, \quad (16)$$

92 with the ground state density for a system of N Fermions given by $n(x) = \sum_{n=1}^N |\phi_n(x)|^2$. Figure 1
 93 illustrates the six lowest in energy eigenstates ($L = 1$) (left panel) and their moduli squared
 (corresponding to single-particle densities) on the right.




Figure 1. The six lowest in energy wave functions (left) and the corresponding single particle densities (right) for the particle-in-a-box model ($L = 1$). For illustrative purposes, all curves are shifted by the corresponding energies, see Eq. (16), in atomic units $\hbar = m = 1$.

94

The second example discussed here is the one-dimensional harmonic oscillator. The potential is given by

$$V(x) = \frac{1}{2} kx^2 = \frac{1}{2} m\omega x^2 \quad \omega = \sqrt{\frac{k}{m}}. \quad (17)$$

We make the usual transformation $y = \sqrt{\alpha}x$ and $\alpha = \frac{m\omega}{\hbar}$. The normalized solutions of the Schrödinger equation with the potential (17) read as

$$\Psi_n(y) = \left(\frac{\alpha}{\pi} \right)^{\frac{1}{4}} \frac{1}{\sqrt{2^n n!}} H_n(y) e^{-\frac{y^2}{2}}, \quad (18)$$

where the $H_n(y)$ are Hermite polynomials, which are solutions of the differential equations $H_n''(y) - 2yH_n'(y) + 2ny = 0$, and n are integer quantum numbers ≥ 0 . The corresponding energies are given by

$$E_n = \left(n + \frac{1}{2} \right) \hbar\omega, \quad n = 0, 1, 2, \dots \quad (19)$$

95 In the following discussion, we set $m = \omega = \hbar = 1$, which corresponds to $\alpha = 1$ and $V(x) = \frac{1}{2}x^2$.
 96 The six lowest in energy single particle wave functions and their corresponding moduli squared are
 97 shown in Figure 2.




Figure 2. The six lowest in energy wave functions (left) and the corresponding single particle densities (right) for the Harmonic oscillator. All curves are shifted by the corresponding energies, see Eq. (19). The quadratic potential is shown in black.

For both examples, we use Eq. (25) to compute the potential. Since the wave functions are given analytically, the potentials can be calculated analytically. We use *Mathematica*[®][22] to accomplish that task. We tested cases up to $N = 6$, and obtained every time the corresponding potential up to a constant. That's not surprising because we have shown in Appendix A that the formalism to calculate those functional derivatives leads to the potential. Equation (30) also obtains a constant that doesn't have a physical meaning. Applying l'Hopital's rule twice at the right boundary for the particles in a box example we obtain the constant:

$$c = \frac{\sum_{n=1}^N \epsilon_n n^2}{\sum_{n=1}^N n^2} = \frac{\pi^2}{2L^2} \frac{\hbar^2 \sum_{n=1}^N n^4}{m \sum_{n=1}^N n^2}. \quad (20)$$

98 Due to the exponential decay of the wave function and the density in the harmonic oscillator example,
 99 only the highest in energy of the occupied orbitals contributes to the constant. We find the constant
 100 to be equal to the eigenvalue of the highest occupied orbital.

101 So far those tests have been performed occupying the lowest N orbitals. We find that any other
 102 combination, mimicking a non-ground state, e.g. by occupying orbitals 1,2,3,4,5 and 7, also leads
 103 to the same potential. The constants for the particles in the box example contains then the sum over
 104 occupied orbitals instead a sum from 1 to N , for the harmonic oscillator the constant is still determined
 105 by the highest occupied orbital.

106 The finding that the formalism yields the potential from the kinetic energy, is not obvious
 107 because the condition $\frac{\delta T_s[n]}{\delta n(\mathbf{r})} = -v_s(\mathbf{r})$ implies the condition of the ground state. On the other hand,
 108 those states correspond to a local extremum so that the expression still holds.

109 Obtaining the potential by a functional differentiation for these examples may sound trivial,
 110 but it demonstrates the power of the procedure and the ability to calculate the functional derivative
 111 analytically. On the other hand, the procedure can be applied to cases where the the single particle
 112 orbitals are given numerically.

So far, we have shown the formalism determines the potential exact (up to a constant) from wave functions by functional differentiation of the kinetic energy, where it is known that all orbitals originate from the same potential, the condition for v -representability. The next step is to determine if a given set of orbitals, forming the density, is v -representable. That can be accomplished by

calculating the potential $v_s(\mathbf{r})$ as shown above, and then checking if each of the orbitals fulfills the Kohn-Sham equation by comparing the right and left-hand sides of the expression,

$$-\frac{1}{2}\nabla_{\mathbf{r}}^2\phi_j(\mathbf{r}) + v_s(\mathbf{r})\phi_j(\mathbf{r}) = \alpha_j\phi_j(\mathbf{r}). \quad (21)$$

113 The determinant is v -representable when the equality holds, (for some α_j), for all orbitals forming the
 114 determinant. It is convenient for the comparison to normalize the left and right-hand side separately
 115 in order to be independent of the α_j .

116 Now we apply this procedure to the case when the density is given. Generally, the task would be
 117 to find the wave function that minimizes the kinetic energy through the procedure of the constrained
 118 search [3]. Cioslowski [16,17] has established the formal procedure for generating the complete set of
 119 antisymmetric wave functions that lead to a given density. For a Slater determinant, the procedure is
 120 outlined in Appendix B. Based on his procedure, for each density, we construct a number of different
 121 Slater determinants that leads to it. For each Slater determinant, we differentiate the expectation value
 122 of kinetic energy with respect to the the density, obtaining $v_s(\mathbf{r})$. Finally, we apply Eq. (21) for every
 123 orbital of the determinant. The results of the test for the case of the particle in the box and harmonic
 124 oscillator are shown in Figures 3 and 4, respectively, for a three particle system.

125 In the figures, the top rows exhibits the density, the second show the set of three mutually
 126 orthonormal orbitals, leading to the density. The third row contains v_s , obtained using Eq. (25), for
 127 each set of orbitals and the remaining rows compare each of the orbitals with the normalized output
 128 corresponding to Eq. (21), where in some the output is multiplied by -1 .

129 In both examples, the target density (top row) is obtained using the three lowest in energy
 130 orbitals (see Eqs. 15 and 18; $L = 1$) for panels A,B and C, the density in panel D is constructed by
 131 using orbitals 2, 3 and 4. We applied Cioslowski's procedure [16] for a single Slater determinant
 132 using auxiliary functions ϕ given in Tables 1 and 2 to obtain new Slater determinants leading to the
 133 given density. Those new orbitals are shown in second row. If the exact orbitals are used, Cioslowski's
 134 procedure returns the same orbitals (column A), otherwise different orbitals are obtained. We note,
 135 the auxiliary functions ϕ do not need to be orthogonal or normalized, as long as they are linearly
 136 independent. The new orbitals, although mutually orthonormal, can have quite strange forms, as
 137 seen in the second row. They do not even have to obey the symmetry of the problem, nor have a
 138 specific number of nodes.

Table 1. initial test functions used in Cioslowski's procedure [16] for the square well example corresponding to Figure 3. The functions in the first row are the exact orbitals, see Eq. (15).

column	test orbital 1	test orbital 2	test orbital 3
A	$\sqrt{2}\sin(\pi x)$	$\sqrt{2}\sin(2\pi x)$	$\sqrt{2}\sin(3\pi x)$
B	1	$\cos(2\pi x)$	$\cos(4\pi x)$
C	1	x	x^2
D	$\frac{1}{4} + x$	$\left(\frac{1}{2} + x\right)^2$	$\left(\frac{3}{4} + x\right)^3$




Figure 3. Results for the square well example, as described in the text. The given density is shown in the top row, where the new orbitals obtained through Cioslowski's procedure [16] using initial functions from Table 1. The potential, obtained using functional differentiation of the expression of the kinetic energy, is plotted in the third row. The three rows at the bottom show the results of the test (see Eq. (21)), where the dashed line corresponds to the left hand side of the equation, and the solid line to the right hand side, equal to the new orbital, both normalized. Some of dashed functions are multiplied by -1 for better comparison, corresponding to a negative α in Eq. (21).




Figure 4. Results analogous to those of Fig. 3, but for the harmonic oscillator example. New orbitals were obtained through Cioslowski's procedure [16] using initial functions from Table 2. The dashed line in subfigure D3 shows $\frac{1}{2}x^2$ for comparison.

Table 2. initial test functions used in Cioslowski's procedure [16] for the harmonic oscillator example corresponding to Figure 4. The functions in the first row are the exact orbitals, see Eq. (18).

column	test orbital 1	test orbital 2	test orbital 3
A	$\frac{1}{\sqrt[4]{\pi}}e^{-\frac{1}{2}x^2}$	$\frac{\sqrt{2}}{\sqrt[4]{\pi}}e^{-\frac{1}{2}x^2}x$	$\frac{1}{\sqrt{2}\sqrt[4]{\pi}}e^{-\frac{1}{2}x^2}(2x^2 - 1)$
B	1	x	x^2
C	1	$\frac{x}{6}$	$\frac{x^2}{18} - 1$
D	1	$\cos\left(\frac{\pi x}{6}\right)$	$\cos\left(\frac{\pi x}{3}\right)$

139 Examining these cases, where different sets of orbitals form the same density (columns B
 140 and C), we see that the left and right hand side of Eq (21) are not the same (see bottom three
 141 rows), even though we know the density to be v -representable. This suggests that most of the
 142 sets generated by Cioslowski's procedure [16] may not correspond to a potential, yet all lead to the
 143 same density by construction. This behavior is expected from the first theorem of Hohenberg and
 144 Kohn: If v -representable, the ground state density determines uniquely the potential and hence the
 145 corresponding ground state wave function.

146 We also applied the procedure to a different density to mimic an excited state (panels D). As seen
 147 on panels D4, D5 and D6, the set of obtained wave functions fail to satisfy Eq (21) and hence are
 148 not v -representable. The choice of the exact orbitals passes the test of v -representability, as already
 149 pointed out above.

150 In order to quantify the deviation of the left and right hand side of Eq. (21) the L^2 norm of the
 151 difference is shown in tables 3 and 4. L^2 norms smaller than the machine epsilon are marked as 0.

Table 3. L^2 norm of the difference of left and right hand side of Eq. (21) for the square well example, corresponding to rows 4–6 in Figure 1.

	A	B	C	D
orbital 1	0	0.5134	0.2508	0.0148
orbital 2	0	0.7795	0.1833	0.1158
orbital 3	0	1.9006	0.1694	0.0428

Table 4. L^2 norm of the difference of left and right hand side of Eq. (21) for the harmonic oscillator example, corresponding to rows 4–6 in Figure 2.

	A	B	C	D
orbital 1	0	0.01390	0.23676	0.33567
orbital 2	0	0	0	0.34787
orbital 3	0	0.00057	0.43090	0.23184

152 3. Conclusions

153 For a non-interacting system, we have shown how v -representability of a given wave function
 154 can be determined. This is accomplished by functional differentiation of the kinetic energy with
 155 respect to the density, and checking whether that derivative is a potential leading to the wave
 156 function. We have shown analytically (in one dimension), that in case of v -representability, the
 157 previously derived expressions for the functional derivative of orbitals lead to the correct result, the
 158 negative of the potential, when applied to terms depending on the diagonal part of the single-particle
 159 density matrix like the expression for the kinetic energy.

160 **Acknowledgments:** Comments by Malcolm Stocks and Don Nicholson are gratefully acknowledged. This work
 161 was supported by by the Critical Materials Institute, an Energy Innovation Hub funded by the U.S. Department

162 of Energy and performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National
 163 Laboratory under Contract DE-AC52-07NA27344.

164 **Author Contributions:** M.D. derived the analytic proof given in Appendix A, all numerical examples and
 165 contributed writing the paper; A.G. contributed in deriving the formalism and writing the paper.

166 **Conflicts of Interest:** The authors declare no conflict of interest.

167 Appendix A. functional differentiation of the kinetic energy (non-interacting)

The non-interacting kinetic energy and its derivative with respect to the density reads (in units $\hbar = m = 1$)

$$T_s = \sum_i \int f_i^*(\mathbf{r}) \left[-\frac{\nabla^2}{2} \right] f_i(\mathbf{r}) \, d\mathbf{r} \quad (22)$$

$$\frac{\delta T_s}{\delta n(\mathbf{r}')} = \sum_i \int \frac{f_i^*(\mathbf{r})}{\delta n(\mathbf{r}')} \left[-\frac{\nabla^2}{2} f_i(\mathbf{r}) \right] \, d\mathbf{r} + \sum_i \int \frac{f_i(\mathbf{r})}{\delta n(\mathbf{r}')} \left[-\frac{\nabla^2}{2} f_i^*(\mathbf{r}) \right] \, d\mathbf{r}, \quad (23)$$

where f_i are solutions of the Kohn-Sham equation, the sum runs over the lowest in energy states. The last expression can be obtained through twice integrating by parts.

For simplicity, we continue in one dimension. The functional derivative of an orbital with respect to the density reads then, using Eq. (13):

$$\begin{aligned} \frac{\delta T_s}{\delta n(x')} &= \sum_i \int \frac{f_i^*(x)}{2n(x)} \left[-\frac{1}{2} f_i''(x) \right] \delta(x-x') \, dx + \sum_i \int \frac{f_i(x)}{2n(x)} \left[-\frac{1}{2} f_i^{*''}(x) \right] \delta(x-x') \, dx \\ &+ \sum_i \int_{x'}^{\infty} \frac{1}{n(x)} \left[f_i^*(x) - f_i^*(x) \frac{n'(x)}{2n(x)} \right] \left[-\frac{1}{2} f_i''(x) \right] \, dx \\ &+ \sum_i \int_{x'}^{\infty} \frac{1}{n(x)} \left[f_i'(x) - f_i(x) \frac{n'(x)}{2n(x)} \right] \left[-\frac{1}{2} f_i^{*''}(x) \right] \, dx \end{aligned} \quad (24)$$

$$\begin{aligned} &= \sum_i \frac{f_i^*(x')}{2n(x')} \left[-\frac{1}{2} f_i''(x') \right] + \sum_i \frac{f_i(x')}{2n(x')} \left[-\frac{1}{2} f_i^{*''}(x') \right] \\ &+ \sum_i \int_{x'}^{\infty} \left[\frac{f_i^*(x)}{n(x)} \left[-\frac{1}{2} f_i''(x) \right] + \frac{f_i'(x)}{n(x)} \left[-\frac{1}{2} f_i^{*''}(x) \right] \right] \, dx \\ &- \sum_i \int_{x'}^{\infty} \left[\frac{f_i^*(x)}{n(x)} \frac{n'(x)}{2n(x)} \left[-\frac{1}{2} f_i''(x) \right] + \frac{f_i(x)}{n(x)} \frac{n'(x)}{2n(x)} \left[-\frac{1}{2} f_i^{*''}(x) \right] \right] \, dx. \end{aligned} \quad (25)$$

If the f_i correspond to the same potential V and are solutions of a Schrödinger (or Kohn-Sham) equation:

$$\left[-\frac{1}{2} f_i''(x) \right] = [\epsilon_i - V(x)] f_i(x) \quad (26)$$

then the expression simplifies to:

$$\begin{aligned}
\frac{\delta T_s}{\delta n(x')} &= \sum_i \epsilon_i \frac{f_i^*(x') f_i(x') + f_i(x') f_i^*(x')}{2n(x')} - \underbrace{\left[\sum_i \frac{f_i^*(x') f_i(x')}{2n(x')} + \sum_i \frac{f_i(x') f_i^*(x')}{2n(x')} \right]}_{=1} V(x') \\
&+ \int_{x'}^{\infty} \left[\frac{\sum_i \epsilon_i (f_i'^*(x) f_i(x) + f_i'(x) f_i^*(x))}{n(x)} \right] dx \\
&- \int_{x'}^{\infty} \underbrace{\left[\frac{\sum_i (f_i'^*(x) f_i(x) + f_i'(x) f_i^*(x))}{n(x)} V(x) \right]}_{=\frac{n'(x)}{n(x)} V(x)} dx \\
&- \int_{x'}^{\infty} \frac{n'(x)}{2n(x)n(x)} \sum_i \epsilon_i [f_i^*(x) f_i(x) + f_i(x) f_i^*(x)] dx \\
&+ \int_{x'}^{\infty} \underbrace{\frac{n'(x)}{2n(x)n(x)} V(x) \sum_i [f_i^*(x) f_i(x) + f_i(x) f_i^*(x)] dx}_{=\frac{n'(x)}{n(x)} V(x)} dx \tag{27}
\end{aligned}$$

$$\frac{\delta T_s}{\delta n(x')} = -V(x') + \frac{\sum_i \epsilon_i f_i^*(x') f_i(x')}{n(x')} + \int_{x'}^{\infty} \left[\frac{\sum_i \epsilon_i f_i^*(x) f_i(x)}{n(x)} \right]' dx \tag{28}$$

$$= -V(x') + \underbrace{\frac{\sum_i \epsilon_i f_i^*(x) f_i(x)}{n(x)} \Big|_{x=\infty}}_{=const} \tag{29}$$

$$\frac{\delta T_s}{\delta n(x')} = -V(x') + c \tag{30}$$

168 For a finite systems the $x = \infty$ in Eq. (29) signifies the right boundary. To derive Eq. (29) integration
169 by parts is used: $\int_a^b f'[x] dx = f[x]|_{x=b} - f[x]|_{x=a}$.

170 Appendix B. Cioslowski's procedure

For a single Slater determinant, we briefly describe how orthogonal orbitals, $\psi(x)$, can be obtained, leading to a given density $n(x) = \sum_{i=1}^N |\psi_i|^2$, using Cioslowski's procedure [16]. From a set linear independent auxiliary functions $\phi(x)$, the $\psi_i(x)$ are given by the expression,

$$\psi_i(x) = \sqrt{f(x)} \sum_{j=1}^N s_{ij} \phi_j(x). \tag{31}$$

The matrix \underline{s} is given by

$$\underline{s} = \underline{S}^{-1/2} \tag{32}$$

with

$$S_{ij} = \int f(x) \phi_i(x) \phi_j(x) dx, \tag{33}$$

and

$$f(x) = n(x) \left[\sum_{ij} \phi_i(x) \left(\underline{S}^{-1} \right)_{ij} \phi_j(x) \right]^{-1}. \tag{34}$$

171 The matrix \underline{S} exists and has to be obtained self-consistently. That can be accomplished by iterating
172 the equations above, e.g. using an identity matrix as starting point.

173 **Bibliography**

- 174 1. Hohenberg, P.; Kohn, W. Inhomogeneous Electron Gas. *Phys. Rev.* **1964**, *136*, B864–B871.
- 175 2. Kohn, W.; Sham, L.J. Self-Consistent Equations Including Exchange and Correlation Effects. *Phys. Rev.*
176 **1965**, *140*, A1133–A1138.
- 177 3. Levy, M. Universal variational functionals of electron densities, first-order density matrices, and natural
178 spin-orbitals and solution of the v -representability problem. *Proceedings of the National Academy of Sciences*
179 **1979**, *76*, 6062–6065.
- 180 4. Levy, M. Electron densities in search of Hamiltonians. *Phys. Rev. A* **1982**, *26*, 1200–1208.
- 181 5. Shimony, A.; Feshbach, H., Eds., Density Functionals for Coulomb Systems. In *Physics as Natural*
182 *Philosophy: Essays in honor of Laszlo Tisza on his 75th Birthday*; Shimony, A.; Feshbach, H., Eds.; M.I.T.
183 Press, 1982; pp. 111–149.
- 184 6. Dreizler, R.; Providencia, J., Eds. *Density Functional Methods in Physics*; Nato ASI Series B123, Plenum
185 Press, 1985.
- 186 7. Kohn, W. v -Representability and Density Functional Theory. *Phys. Rev. Lett.* **1983**, *51*, 1596–1598.
- 187 8. Englisch, H.; Englisch, R. Hohenberg-Kohn theorem and non- V -representable densities. *Physica A:*
188 *Statistical Mechanics and its Applications* **1983**, *121*, 253 – 268.
- 189 9. Chen, J.; Stott, M.J. v -representability for noninteracting systems. *Phys. Rev. A* **1993**, *47*, 153–160.
- 190 10. Parr, R.G.; Yang, C.Y. *Density Functional Theory of Atoms and Molecules*; Oxford University Press, Oxford,
191 1989.
- 192 11. Dreitzler, R.M.; Gross, E.K.U. *Density Functional Theory*; Springer Verlag, Berlin, New York, 1990.
- 193 12. Gonis, A.; Zhang, X.G.; Däne, M.; Stocks, G.; Nicholson, D. Reformulation of density functional theory
194 for N -representable densities and the resolution of the v -representability problem. *Journal of Physics and*
195 *Chemistry of Solids* **2016**, *89*, 23–31.
- 196 13. Gonis, A.; Däne, M.; Nicholson, D.; Stocks, G. Computationally simple, analytic, closed form solution of
197 the Coulomb self-interaction problem in Kohn-Sham density functional theory. *Solid State Communications*
198 **2012**, *152*, 771 – 774.
- 199 14. Däne, M.; Gonis, A.; Nicholson, D.; Stocks, G. On a solution of the self-interaction problem in Kohn-Sham
200 density functional theory. *Journal of Physics and Chemistry of Solids* **2014**, *75*, 1160 – 1178.
- 201 15. Däne, M.; Gonis, A.; Nicholson, D.; Stocks, G. Solving the self-interaction problem in Kohn-Sham density
202 functional theory: Application to atoms. *Journal of Physics and Chemistry of Solids* **2015**, *79*, 55 – 65.
- 203 16. Cioslowski, J. Density Functionals for the Energy of Electronic Systems: Explicit Variational Construction.
204 *Phys. Rev. Lett.* **1988**, *60*, 2141–2143.
- 205 17. Cioslowski, J. Density driven self-consistent field method. II. construction of all one-particle
206 wavefunctions that are orthonormal and sum up to a given density. *International Journal of Quantum*
207 *Chemistry* **1989**, *36*, 255–262.
- 208 18. Lieb, E.H. Density Functionals for Coulomb Systems. In *Physics as Natural Philosophy, Essays in Honor of*
209 *Laszlo Tisza*; Shimony, A.; Feshbach, H., Eds.; MIT Press Cambridge, Massachusetts, 1982.
- 210 19. Macke, W. Zur wellenmechanischen Behandlung von Vielkörperproblemen. *Annalen der Physik* **1955**,
211 *452*, 1–9. 6. Folge, Band 17, Heft 1 (in German).
- 212 20. Harriman, J.E. Orthonormal orbitals for the representation of an arbitrary density. *Phys. Rev. A* **1981**,
213 *24*, 680–682.
- 214 21. Zumbach, G.; Maschke, K. New approach to the calculation of density functionals. *Phys. Rev. A* **1983**,
215 *28*, 544–554.
- 216 22. Mathematica Version 10.4; Wolfram Research, Inc.: Champaign, IL, USA, 2016.